

Product Information

**IgG2b-FITC Isotype Control from murine myeloma
Clone MOPC-141**
purified immunoglobulin

Catalog Number **F6647**

Product Description

The MOPC-141 tumor line that produces mouse IgG2b,κ, is a mineral oil induced plasmacytoma. The product is originated and carried intraperitoneally in BALB/c mice.

Identity and purity of the immunoglobulin is established by immunoelectrophoresis (IEP) and Ouchterlony Double Diffusion (ODD), prior to conjugation. Electrophoresis of the purified immunoglobulin, followed by diffusion against anti-mouse whole serum and anti-mouse IgG2b results in single arcs of precipitation. By ODD the purified immunoglobulin preparation is non-reactive with antisera to mouse IgA, IgM, IgG1, IgG2a, and IgG3.

The specificity of staining by monoclonal antibodies to human CD antigens should be verified by establishing the amount of non-specific binding to the target cell population. It is recommended that a non-reactive immunoglobulin of the same isotype, concentration and F/P molar be included as a negative control for each monoclonal antibody reagent used in flow cytometry or other immunoassays.

Reagent

The IgG2b,κ, purified from ascites fluid, is conjugated to fluorescein isothiocyanate and provided as a solution in 0.01 M PBS, pH 7.4, containing 1.0% BSA and 15 mM sodium azide as a preservative.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C. For extended storage, the solution may be frozen in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

Note: Store product protected from light.

Product Profile

Equal concentrations of isotype control and primary antibody are recommended for use in flow cytometry.

Reactivity

When evaluated in flow cytometry, the product did not stain human peripheral blood lymphocytes (PBLs), obtained from whole blood lysis.

NRC,KAA,PHC 09/08-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.