

**MONOCLONAL Anti-Human CD7 Clone 3A1
FITC Conjugate
Purified Mouse Immunoglobulin**Product No. **F0647**

Monoclonal Anti-Human CD7 (mouse IgG2b isotype) is derived from the hybridoma produced by the fusion of mouse myeloma cell line NS-1 and splenocytes from Balb/c mice immunized with the human T cell line HSB-2. The isotype is determined using Sigma Immunotype™ Kit (Sigma Stock No. ISO-1) and by a double diffusion assay using Mouse Monoclonal Antibody Isotyping Reagents (Sigma Stock No. ISO-2). The product is prepared by conjugation of fluorescein isothiocyanate (FITC) Isomer I to purified CD7 monoclonal antibody. The conjugate is then purified by gel filtration to remove unbound FITC, no free FITC is detectable. The conjugate is provided (200 µg/ml) in 0.01 M phosphate buffered saline, pH 7.4, containing 1% BSA with 0.1% sodium azide (see MSDS)* as a preservative.

Description

Monoclonal Anti-Human CD7 recognizes the CD7 human lymphocyte surface antigen glycoprotein (molecular weight 40kD). This antigen is present on more than 85% of the normal peripheral blood lymphocytes and most NK cells. It is expressed by a majority of thymocytes, T-ALL, and T-cell lymphoblastic lymphomas. Monoclonal Anti-Human CD7 does not stain most B lymphocytes or granulocytes. Blocking studies suggest that there is identity between this molecule and the receptor for IgM (FcµR). The epitope recognized by this clone is sensitive to routine formalin fixation and paraffin embedding.

Performance

When assayed by flow cytometric analysis, using 10 µl of the antibody to stain 1 x 10⁶ cells, a fluorescence intensity is observed similar to that obtained with saturating monoclonal antibody levels. The percent population positive is also at the maximum percentage positive using saturating monoclonal antibody levels.

F/P Molar Ratio: 3-8**Uses**

Monoclonal Anti-Human CD7, FITC conjugate may be used for:

1. Analysis of T cell and NK subsets.
2. Classification of acute leukemias and lymphomas.
3. Characterization of immunodeficiencies.

Storage

Store at 2-8°C. Protect from prolonged exposure to light. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

* Due to the sodium azide content, a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazardous and safe handling practices.

Note: In order to obtain best results in different preparations, it is recommended that each individual user determine their optimum working dilutions by titration assay.

Procedure for Direct Immunofluorescent Staining
Reagents and Materials Needed but Not Supplied

1. a. Whole human blood collected by standard clinical blood evacuation tubes with EDTA, ACD-A or heparin anticoagulant **or**
b. Human cell suspension (e.g., peripheral blood mononuclear cells isolated on HISTOPAQUE® (Sigma Product No. 1077-1)).
2. Diluent: 0.01M phosphate buffered saline (PBS), pH 7.4, containing 1% BSA and 0.1% NaN₃.
3. FITC conjugated, isotype-matched, non-specific mouse immunoglobulin (negative control, Sigma Product No. F6647).
4. 12 x 75 mm test tubes.
5. Adjustable micropipet.
6. Centrifuge.
7. Counting chamber.
8. Trypan blue (Sigma Product No. T0776), 0.2% in 0.01 M PBS, pH 7.4.
9. 2% paraformaldehyde in PBS.
10. Whole blood lysing solution.
11. Flow cytometer.

Procedure

1. a. Use 100 μ l of whole blood **or**
 - b. Adjust cell suspension to 1×10^7 cells/ml in diluent. Cells should be >90% viable as determined by dye exclusion (e.g., trypan blue). For each sample, add 100 μ l or 1×10^6 cells per tube.
2. Add 10 μ l of conjugate to tube(s) containing cells to be stained. Vortex tube gently. Incubate the cells at room temperature (18 - 22°C) for 30 minutes. Proper controls to be included for each sample are:
 - a. An autofluorescence control: 10 μ l diluent in place of monoclonal antibody, followed by steps 3 - 7.
 - b. A negative staining control: 10 μ l of FITC conjugated, isotype-matched non-specific mouse immunoglobulin (Sigma Product No. F6647) at the same concentration as test antibody followed by steps 3 - 7.
3. a. If whole blood is used, use lysing solution after incubation and wash cells according to manufacturer's instructions.
 - b. If a mononuclear cell suspension is used, proceed to Step. 4.
4. Add 2 ml of diluent to all tubes.
5. Pellet cells by centrifugation at 500 x G for 10 minutes.
6. Remove supernatant by careful aspiration.
7. Resuspend cells in 0.5 ml of 2% paraformaldehyde. Analyze in a flow cytometer according to manufacturer's instructions.

Quality Control

It is advisable to run the appropriate negative controls. Negative controls establish background fluorescence and non-specific binding of the primary and secondary antibodies. The ideal negative control reagent is a mouse monoclonal or myeloma protein which has no reactivity with human cells. It should be isotype-matched to the antibody and of the same concentration and F/P molar ratio as the antibody. The degree of autofluorescence or negative control reagent fluorescence will vary with the type of cells under study and the sensitivity of the instrument used.

References

- Haynes, B.F., et al., Proc. Nat. Acad. Sci. USA, **76**, 5829 (1979).
Haynes, B.F., et al., Proc. Natl. Acad. Sci. USA, **77**, 2914 (1980).
Leucocyte Typing IV, Knapp, W., et al., (Eds.), Oxford Press, New York, 1075 (1989).
Leucocyte Typing IV, Knapp, W., et al., (Eds.), Oxford Press, New York, 1076 (1989).
Eisenbarth, G.S., et al., J. Immunol., **124**, 1237 (1980).
Sutherland, D.R., et al., J. Immunol., **133**, 327 (1984).
Aruffo, A. and B. Seed, EMBO J., **6**, 3313 (1987).

1/98

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications.

Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply.

Please see reverse side of the invoice or packing slip.