

Product Information

Monoclonal Anti-Interleukin-1 α , clone 3405
produced in mouse, purified immunoglobulin

Catalog Number **I7659**

Product Description

Monoclonal Anti-Interleukin-1 α (IL-1 α) (mouse IgG1 isotype) is purified from a hybridoma produced by the fusion of mouse myeloma cells and B cells from a mouse immunized with recombinant human Interleukin-1 α (GeneID 3552) expressed and purified from *Escherichia coli*. The antibody is purified by Protein G affinity chromatography.

Monoclonal Anti-Interleukin-1 α recognizes human Interleukin-1 α . Applications include flow cytometry, and neutralization.

Interleukin-1 (IL-1) is a name that designates two proteins, IL-1 α and IL-1 β , which are the products of distinct genes, but which share approximately 25% amino acid sequence identity. Both bind to the same cell surface receptor, and elicit nearly identical biological responses. IL-1 α is synthesized as a precursor protein that lacks a signal peptide. IL-1 α precursor is localized to the nucleus, cytosol, and plasma membrane. Mature IL-1 α is generated via cleavage by the cysteine protease calpain. A small percentage of total cellular IL-1 α precursor can be found on the surface of various cells. This membrane bound IL-1 α is probably a glycosylated or myristoylated form of the cytokine.

Interleukin-1 (IL-1), originally known as lymphocyte activating factor (LAF), activates T cells and lymphocytes, which then proliferate and secrete interleukin-2.¹ IL-1 is primarily released from stimulated macrophages and monocytes, but also is released from several other cell types,² and is thought to play a key role in inflammatory and immune responses.³ Other synonyms for IL-1 include: endogenous pyrogen (EP), mitogenic protein (MP), helper peak-1 (HP-1), T cell replacing factor III (TRF III or TRFH), B cell activating factor (BAF) and B cell differentiation factor (BDF).⁴

Reagent

Supplied lyophilized from a 0.2 μ m filtered solution of phosphate buffered saline with 5% trehalose.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

To one vial of lyophilized powder, add 1 mL of 0.2 μ m filtered PBS to produce a 0.5 mg/mL stock solution. If aseptic technique is used, no further filtration should be necessary for use in cell culture environments.

Storage/Stability

Prior to reconstitution, store at -20°C . Reconstituted product may be stored at $2-8^{\circ}\text{C}$ for up to one month. For extended storage, freeze in working aliquots at -20°C . Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended.

Neutralization

To measure the ability of the antibody to neutralize the bioactivity of recombinant human Interleukin-1 α on D10.G4.1 cells, recombinant human Interleukin-1 α was incubated with various concentrations of the antibody for 1 hour at 37°C in a 96 well plate. Following this preincubation period, D10.G4.1 cells were added. The assay mixture in a total volume of 200 μ L, containing antibody at the concentrations indicated, recombinant human Interleukin-1 α at 50 pg/mL, Concanavalin A at 1.25 μ g/mL and cells at 5×10^4 cells/mL, was incubated at 37°C for 72 hours in a humidified CO_2 incubator. ^3H -thymidine was added during the final 4 hours of incubation. The cells were subsequently harvested onto glass fiber filters and the ^3H -thymidine incorporated into DNA was determined.

Human Interleukin-1 α stimulates ^3H -thymidine incorporation by murine T-helper D10.G4.1 cells in a dose-dependent manner. The ED_{50} for this effect is typically 3-7 pg/mL.⁵

The Neutralization Dose₅₀ (ND₅₀) for this antibody is defined as that concentration of antibody required to yield one-half maximal inhibition of the cytokine activity on a responsive cell line, when that cytokine is present at a concentration just high enough to elicit a maximum response.

Product Profile

Flow Cytometry: this antibody can be used to detect membrane-bound forms of human Interleukin-1 α . Dilute this antibody to ~25 $\mu\text{g}/\text{mL}$ and add 10 μL of this solution to $1\text{-}2.5 \times 10^5$ cells in a total reaction volume not exceeding 200 μL . The binding of unlabeled monoclonal antibodies may be visualized by using a secondary developing reagent such as goat anti-mouse IgG conjugated to a fluorochrome.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

Endotoxin: $<0.1 \text{ EU}/\mu\text{g}$ antibody as determined by the LAL method.

References

1. Gery, I., et al., *J. Exp. Med.*, **136**, 128 (1972).
2. Oppenheim, J., et al., *Immunol. Today*, **7**, 45 (1986).
3. Durum, S., et al., *Ann. Rev. Immunol.*, **3**, 263 (1985).
4. Aarden, L., et al., *J. Immunol.*, **123**, 2928 (1979).
5. Symons, J. A., et al., in *Lymphokines and Interferons, a practical approach*, IRL Press, M.J. Clemens, A. G. Morris and A.J.H. Gearing, eds. p. 272 (1987).

RC,PHC 07/11