

Product Information

Phosphatase, Alkaline from *Escherichia coli*

Lyophilized powder, 30-60 units/mg protein (in glycine buffer)

P5931

CAS Registry Number: 9001-78-9

Enzyme Commission (EC) Number: 3.1.3.1

Synonyms: Orthophosphoric-monoester phosphohydrolase (alkaline optimum), alkaline phosphomonoesterase; phosphomonoesterase; glycerophosphatase; alkaline phosphohydrolase; alkaline phenyl phosphatase

Storage Temperature: -20 °C

Molecular Mass: ~94 kDa¹

Product Description

Alkaline phosphatase is a dimeric metalloprotein, containing zinc and magnesium, that catalyzes the hydrolysis and the transphosphorylation of various phosphomonoesters. In *E. coli* specifically, alkaline phosphatase is normally located in the periplasm.¹⁻¹⁴ Though the subunits are believed to be coded by the same gene, the subunits develop molecular heterogeneity after translation.

The enzyme belongs to a group of phosphohydrolases that have an optimal pH for *in vitro* activity of ~10. The actual optimum varies, depending on the nature and concentration of the substrate, the type of buffer, the phosphate acceptor, and to some extent the nature of the isoenzymes. As the enzyme generally requires divalent ions (Mg²⁺, Co²⁺, or Mn²⁺) as an activator and contains Zn²⁺ as a constituent metal ion, chelating agents and inorganic phosphates inhibit the enzyme.

Several dissertations have cited use of this product in their research.^{15,16}

Precautions and Disclaimer

This product is for R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Product

The product is supplied as a lyophilized powder containing Tris buffer salts, MgSO₄, and ZnSO₄.

Unit Definition: One unit will hydrolyze 1 μmole of 4-nitrophenyl phosphate per minute at pH 10.4 at 37 °C (in glycine buffer).

The enzyme activity is determined by measuring the increase in absorbance at 410 nm that results from the hydrolysis of *p*-nitrophenylphosphate to *p*-nitrophenol. The reaction is performed in 87 mM glycine buffer with 0.9 mM MgCl₂, 0.87 mM ZnCl₂, and 6 mM *p*-nitrophenyl phosphate at pH 10.4 at 37 °C. Immediately before use, the enzyme solution is prepared at 0.1-0.2 unit/mL of cold 1.0 mM MgCl₂.

Preparation Instructions

The product is soluble in aqueous solution with 1 mM MgCl₂ (1 mg/mL), yielding a clear, colorless solution.

Storage/Stability

The product has a recommended rested date of 4 years when unopened and properly at -20 °C. This product may be stored under desiccator conditions.

References

1. Bučević-Popović, V. *et al.*, *Biochimie*, 86(6), 403-409 (2004).
2. Plocke, D. J. *et al.*, *Biochemistry*, 1(3), 373-378 (1962).
3. Simpson, R. T. *et al.*, *Biochemistry*, 7(12), 4336-4342 (1968).
4. Coleman, J. E., *Annu. Rev. Biophys. Biomol. Struct.*, 21, 441-483 (1992).
5. Sambrook, J. *et al.*, in *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press (Cold Spring Harbor, NY), p. 5.72 (1989).
6. Mossner, E., *et al.*, *Hoppe Seyler's Z. Physiol. Chem.*, 361(4), 543-549 (1980).
7. Reid, T. W., and Wilson, I. B., *The Enzymes*, 4, 373-415 (1971).
8. Anderson, R. A., *et al.*, *Proc. Nat. Acad. Sci. USA*, 72(8), 2989-2993 (1975).
9. Malamy, M. H., and Horecker, B. L., *Biochemistry*, 3(12), 1893-1897 (1964).
10. Stadtman, T. C., "Alkaline Phosphatases", in *The Enzymes*, 2nd ed., Vol. 5 (Boyer, P. D. *et al.*, eds.). Academic Press (New York, NY), pp. 55-71 (1961).
11. Bloch, W., and Gorby, M. S., *Biochemistry*, 19(22), 5008-5018 (1980).
12. Kim, E. E., and Wyckoff, H. W., *J. Mol. Biol.*, 218(2), 449-464 (1991).
13. Xu, X., and Kantrowitz, E. R., *Biochemistry*, 30(31), 7789-7796 (1991).
14. Zhang, L. *et al.*, *Biochem. J.*, 392(Pt 2), 407-415 (2005).
15. Jalland, Clara Maria Osnes, "Influence of base-excision-repair pathway enzymes on prion pathogenesis". Norwegian University of Life Sciences, Ph.D. dissertation, pp. 89, 98 (2017).
16. Verma, Nipun, "TET proteins safeguard bivalent promoters from de novo methylation in hESCs". Weill Cornell Graduate School of Medical Sciences, Ph.D. dissertation, p. 121 (2017).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2002-2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

P5931pis Rev 12/23 AB, MP

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

Terms and Conditions of Use

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

**MILLIPORE
SIGMA**