

Product Information

**PDGFR α (D842V), active, GST-tagged, human
PRECISIO® Kinase
recombinant, expressed in *Sf9* cells**

Catalog Number **SRP5303**
Storage Temperature -70°C

Synonyms: CD140A, PDGFR2, MGC74795,
Rhe-PDGFR α

Product Description

PDGFR α (platelet-derived growth factor receptor α) is a member of the PDGFR family of membrane receptors with intrinsic tyrosine kinase activity. Aberrant expression of PDGFR α has been linked to developmental abnormalities in vertebrate models and has been implicated in multiple disease states in humans. There is widespread expression of PDGFR α in renal cell types involved in fibrotic and sclerosing processes.¹ PDGF and its receptor PDGFR α are inducers of fibrosis in the repair phase of inflammatory bowel disease and they may also be involved in the active inflammatory phase.² PDGFR α (D842V) is one of the mutant forms of PDGFR α .

Recombinant human PDGFR α (D842V) (550-end) was expressed by baculovirus in *Sf9* insect cells using an N-terminal GST-tag. The gene accession number is NM_006206. It is supplied in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, and 25% glycerol.

Molecular mass: ~95 kDa

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

The product ships on dry ice and storage at -70°C is recommended. After opening, aliquot into smaller quantities and store at -70°C . Avoid repeated handling and multiple freeze/thaw cycles.

Figure 1.
SDS-PAGE Gel of Typical Lot:
 $\geq 70\%$ (SDS-PAGE, densitometry)





Figure 2.
Specific Activity of Typical Lot:
43–65 nmole/min/mg



Procedure

Preparation Instructions

Kinase Assay Buffer – 25 mM MOPS, pH 7.2, 12.5 mM glycerol 2-phosphate, 20 mM MgCl_2 , 25 mM MnCl_2 , 5 mM EGTA, and 2 mM EDTA. Just prior to use, add DTT to a final concentration of 0.25 mM.

Kinase Dilution Buffer – Dilute the Kinase Assay Buffer 5-fold with a 50 ng/ μL BSA solution.

Kinase Solution – Dilute the active PDGFR α (D842V) (0.1 $\mu\text{g}/\mu\text{L}$) with Kinase Dilution Buffer to the desired concentration.

Note: The specific activity plot may be used as a guideline (see Figure 2). It is recommended the researcher perform a serial dilution of active PDGFR α (D842V) kinase for optimal results.

10 mM ATP Stock Solution – Dissolve 55 mg of ATP in 10 mL of Kinase Assay Buffer. Store in 200 μL aliquots at -20°C .

γ - ^{33}P -ATP Assay Cocktail (250 μM) – Combine 5.75 mL of Kinase Assay Buffer, 150 μL of 10 mM ATP Stock Solution, 100 μL of γ - ^{33}P -ATP (1 mCi/100 μL). Store in 1 mL aliquots at -20°C .

Substrate Solution – MBP Protein substrate diluted in distilled water to a final concentration of 1 mg/mL.

1% phosphoric acid solution – Dilute 10 mL of concentrated phosphoric acid to a final volume of 1 L with water.

Kinase Assay

This assay involves the use of the ^{33}P radioisotope. All institutional guidelines regarding the use of radioisotopes should be followed.

1. Thaw the active PDGFR α (D842V), Kinase Assay Buffer, Substrate Solution, and Kinase Dilution Buffer on ice. The γ - ^{33}P -ATP Assay Cocktail may be thawed at room temperature.
2. In a pre-cooled microcentrifuge tube, add the following solutions to a volume of 20 μL :
 - 10 μL of Kinase Solution
 - 5 μL of Substrate Solution
 - 5 μL of cold water (4°C)
3. Set up a blank control as outlined in step 2, substituting 5 μL of cold water (4°C) for the Substrate Solution.
4. Initiate each reaction with the addition of 5 μL of the γ - ^{33}P -ATP Assay Cocktail, bringing the final reaction volume to 25 μL . Incubate the mixture in a water bath at 30°C for 15 minutes.
5. After the 15 minute incubation, stop the reaction by spotting 20 μL of the reaction mixture onto an individually pre-cut strip of phosphocellulose P81 paper.

6. Air dry the pre-cut P81 strip and sequentially wash in the 1% phosphoric acid solution with constant gentle stirring. It is recommended the strips be washed a total of 3 times of ~ 10 minutes each.
7. Set up a radioactive control to measure the total γ - ^{33}P -ATP counts introduced into the reaction. Spot 5 μL of the γ - ^{33}P -ATP Assay Cocktail on a pre-cut P81 strip. Dry the sample for 2 minutes and read the counts. Do not wash this sample.
8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.
9. Determine the corrected cpm by subtracting the blank control value (see step 3) from each sample and calculate the kinase specific activity

Calculations:

1. Specific Radioactivity (SR) of ATP (cpm/nmole)

$$\text{SR} = \frac{\text{cpm of } 5 \mu\text{L of } \gamma\text{-}^{33}\text{P-ATP Assay Cocktail}}{\text{nmole of ATP}}$$

cpm – value from control (step 7)
nmole – 1.25 nmole (5 μL of 250 μM ATP Assay Cocktail)

2. Specific Kinase Activity (SA) (nmole/min/mg)

$$\text{nmole/min/mg} = \frac{\Delta\text{cpm} \times (25/20)}{\text{SR} \times \text{E} \times \text{T}}$$

SR = specific radioactivity of the ATP (cpm/nmole ATP)

Δcpm = cpm of the sample – cpm of the blank (step 3)

25 = total reaction volume

20 = spot volume

T = reaction time (minutes)

E = amount of enzyme (mg)

References

1. Floege, J. et al., Expression of PDGF alpha-receptor in renal arteriosclerosis and rejecting renal transplants. J. Am. Soc. Nephrol., **9**(2), 211-23 (1998).
2. Kumagai, S. et al., Platelet-derived growth factor and its receptors are expressed in areas of both active inflammation and active fibrosis in inflammatory bowel disease. Tohoku J. Exp. Med., **195**(1), 21-33 (2001).

PRECISIO is a registered trademark of Sigma-Aldrich Co. LLC.

RC,MAM 12/12-1