

Product Information

Anti-Staphylococcal Enterotoxin B Antibody, Mouse Monoclonal

Clone SEB4, Purified from Hybridoma Cell Culture

SAB4200859

Product Description

Monoclonal Anti- Staphylococcal Enterotoxin B antibody (mouse IgG1 isotype) is derived from the SEB4 hybridoma, produced by the fusion of mouse myeloma cells and splenocytes from a mouse immunized with staphylococcal enterotoxin B from Staphylococcus aureus as immunogen. The isotype is determined by ELISA using Mouse Monoclonal Antibody Isotyping Reagents (Sigma ISO-2). The antibody is purified from culture supernatant of hybridoma cells.

Monoclonal Anti-Staphylococcal Enterotoxin B antibody specifically recognizes staphylococcal enterotoxin B (SEB) and has no cross reactivity with staphylococcal enterotoxin A (SEA), cholera toxin, or pseudomonas exotoxin A. The antibody may be used in various immunochemical techniques including Immunoblotting and ELISA. Detection of the staphylococcal enterotoxin B bands by Immunoblotting is specifically inhibited by the immunogen.

Staphylococcus aureus is a Gram-positive bacterium, that causes disease in the human population including food poisoning and toxic shock syndrome. 1 S. aureus expresses several secreted virulence factors, such as various enzymes, cytotoxins, exotoxins, and exfoliative toxins.1 Exotoxins include more than 20 serologically classified staphylococcal enterotoxins (SEs). The best characterized are the SEs A, B through V, and toxic shock syndrome toxin-1 (TSST-1). These enterotoxins are similar in activity, sequence, structure, and molecular mass (25-30 kDa).1-3 SEA and SEB are known as superantigens due to their ability to bind class II MHC molecules on antigen presenting cells. This binding stimulates extensive T-cell activation followed by massive cytokine release leading to an acute toxic shock.¹⁻⁴ SEB is also classified as a category B select agent and is important toxicant in biodefense research.5-9

SE proteins have a significant resistance to heat and acid, and are also resistant to gastrointestinal proteases including pepsin, trypsin, rennin, and papain. Thus, inactivating the *S. aureus* bacteria may not be sufficient to eliminate the risk of its superantigens from causing food poisoning.¹

SEB contamination can be caused by both its ingestion (resulting with food poisoning) or by inhalation (resulting with respiratory symptoms of cough and dyspnea), therefore SEB detection is a major challenge in the food industry. $^{1,3,5-10}$

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody Concentration: ~ 1.0 mg/mL

Precautions and Disclaimer

For research use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

1

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.



Product Profile

Immunoblotting: a working concentration of 0.5-1.0 µg/mL is recommended using staphylococcal enterotoxin B.

ELISA: a working concentration of 0.12-0.25 μg/mL is recommended using 1 μg/mL staphylococcal enterotoxin B for coating.

Note: In order to obtain best results in different techniques and preparations it is recommended to determine optimal working concentration by titration test.

References

- Pinchuk IV., et al., Toxins (Basel), 2, 2177-97 (2010).
- Bergdoll MS., Enterotoxins. In: Staphylococci and Staphylococcal Infections, Easmon CSF. and Adlam C. (eds), Academic Press, London 559-598 (1983).
- 3. Miron N. and Miron MM., *Microbiol Immunol.*, **54**, 769-77 (2010).
- Pontzer CH., et al., Proc Natl Acad Sci U S A., 88, 125-8 (1991).
- Alefantis T., et al., Mol Cell Probes, 18, 379-82 (2004).
- Karauzum H., et al., J Biol Chem., 287, 25203-15 (2012).
- 7. Pita R. and Romero A., Forensic Sci Rev., **26**, 85-96 (2014).
- Rusnak JM., et al., Emerg Infect Dis., 10, 1544-9 (2004).
- Rajagopalan G., et al., Shock., 25, 647-56 (2006).
- 10. Konry T., et al., *Biosens Bioelectron.*, **22**, 2230-6 (2007).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.



2