



3050 Spruce Street  
Saint Louis, Missouri 63103 USA  
Telephone (800) 325-5832 (314) 771-5765  
Fax (314) 286-7828  
email: [techserv@sial.com](mailto:techserv@sial.com)  
[sigma-aldrich.com](http://sigma-aldrich.com)

## Product Information

### RIBONUCLEOSIDE VANADYL COMPLEXES, 200 mM

Product No. R 3380

Lot 045K6126

Store below -20 °C

#### Product Summary

A 20 mM solution inhibits 0.00025 units/ml of RNase A.

#### Comments

**Note:** SDS and EDTA or other chelating agents may dissociate and inactivate ribonucleoside vanadyl complexes.

#### Introduction

Ribonucleoside vanadyl complexes (RVC) have been used as ribonuclease inhibitor during cell lysis and cDNA production by reverse transcriptase. DNase I is not inhibited by 20 mM RVC and consequently DNA can be degraded with DNase I while using RVC to protect RNA from contaminating ribonucleases. RVC inhibit most nucleases with the exception of S1 nuclease, DNase I, and *Bacillus cereus* ribonuclease. RVC are not compatible with *in vitro* translation systems, but are tolerated when included with mRNA microinjected into frog oocytes.

Phenol extraction can be used to remove RVC from samples. If 8-hydroxyquinoline is included as an antioxidant with phenol, removal of RVC by successive phenol extraction is easily monitored. The orange colored phenol solution turns black as RVC are removed, but remains orange when all RVC have been extracted from the aqueous phase.

#### Suitability

Ribonucleoside vanadyl complexes at 20 mM final concentration, 50 mM Tris-HCl, pH 8.0 and 2 µg of tRNA were incubated with 0.0001-0.003 Kunitz units/ml of RNase A for 30 minutes at 37 °C in a 50 µl reaction. An aliquot of this reaction was analyzed by electrophoresis on a 10-20% linear gradient polyacrylamide gel. Ribonucleoside vanadyl complexes were found to inhibit approx. 0.0002 Kunitz units/ml of RNase A. Detection limit: Degradation by  $3 \times 10^6$  Kunitz units of RNase A detectable.

#### References:

1. Berger, S. L., and Birkenmeier, C. S., *Biochemistry*, **18**, 5143 (1979).
2. Puskas, R. S., et al., *Biochemistry*, **21**, 4602 (1982).

9/05

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply.

Please see reverse side of the invoice or packing slip.