3050 Spruce Street, Saint Louis, MO 63103 USA Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757 email: techservice@sial.com sigma-aldrich.com

# Anti-Mucolipin-2 (C-terminal)

produced in rabbit, affinity isolated antibody

Catalog Number M1196

## **Product Description**

Anti-Mucolipin-2 (C-terminal) is produced in rabbit using as immunogen a synthetic peptide corresponding to amino acids 534-549 of human mucolipin-2 (GeneID: 255231), conjugated to KLH. This sequence has low homology with rat (73% identity) and mouse (66% identity) mucolipin-2. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-Mucolipin-2 (C-terminal) specifically recognizes human mucolipin-2 by immunoblotting (~60 kDa). Additional bands may be observed at ~120 kDa and ~35 kDa, due to aggregation and degradation of mucolipin-2, respectively. Staining of the mucolipin-2 band is specifically inhibited by the immunizing peptide.

Mucolipidosis type IV (MLIV) is an autosomal recessive, neurodegenerative disorder caused by mutations in the MCOLN1 gene that encodes mucolipin-1 (also termed TRP-ML1, MLN1, ML1 mucolipidin). 1, 2 MLVI is associated with severe psychomotor retardation and ophthalmologic defects. MLIV is a lysosomal storage disorder associated with lysosomal accumulation of sphingolipids, phospholipids and mucopolysaccharides. Unlike other mucolipidoses, lysosomal hydrolase activity is not impaired in MLVI. Rather, MLVI pathophysiology has been linked to deficiency in membrane trafficking and organelle dynamics in the late endocytic pathway.<sup>2</sup> MLN1 shares significant sequence homology with the TRP superfamily of cation channels, characterized by permeability to monovalent cations and Ca2+. MLN1 is thought to function as a proton-leak channel in lysosomes, regulating lysosomal pH and hydrolytic activity. MLN1 has been localized to late endosomes and lysosomes. In addition to MLN1, mammals encode two other highly related proteins, MLN2/TRPML2 and MLN3/TRPML3. Mutations in mouse mucolipin-3 (MLN3, TRPML3) encoded by the MCOLN3 gene, are associated with deafness and pigmentation defects in varitint-waddler mice.4 TRPMLs have been shown to interact to form homoand heteromultimers resulting in a complex pattern of localization in both the lysosomal and endoplasmic reticulum compartments.5

### Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~1.0 mg/mL

#### **Precautions and Disclaimer**

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

## Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

## **Product Profile**

 $\frac{Immunoblotting}{0.3\text{-}0.6~\mu\text{g/mL}} \text{ a working antibody concentration of } 0.3\text{-}0.6~\mu\text{g/mL} \text{ is recommended using HEK-293T cells expressing human mucolipin-2}.$ 

**Note**: In order to obtain the best results and assay sensitivity in various techniques and preparations, we recommend determining optimal working concentrations by titration.

## References

- Bach, G., et al., Eur. J. Physiol., 451, 313-317 (2005).
- 2. Bargal, R., et al., *Nature Genet.*, **26**, 118-123 (2000).
- 3. Soyombo, A.A., et al., *J. Biol. Chem.*, **281**, 7294-7301 (2006).
- Di Palma, F., et al., Proc. Natl. Acad. Sci. USA, 99, 14994-14999 (2002).
- 5. Zeevi, D.A., et al., *Biochim. Biophys. Acta,* **1772**, 851-858 (2007).

ER,KAA,PHC 12/07-1

Sigma brand products are sold through Sigma-Aldrich, Inc.
Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications.
Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply.

SIGMA-ALDRICH™