Technical Data Sheet

DCA (Differential Clostridial Agar) acc. to WEENK et al.

For the enumeration of sulfite-reducing clostridia in dried foods.

1102590500

Mode of Action

The DCA medium consists of a nutritionally rich base medium, including starch to promote spore germination. Resazurin is added as a redox indicator, turning red at high redox potential, indicating aerobic conditions. Sulfite and an iron source are added as indicators. Sulfite-reducing clostridia produces sulfide from sulfite, which gives a black precipitate with the iron present in the medium. Sulfite-reducing clostridia are enumerated as black colonies.

Typical Composition (g/L)

Peptone from casein	5.0
Peptone from meat	5.0
Meat extract	8.0
Starch	1.0
D(+)glucose	1.0
Yeast extract	1.0
Cysteinium chloride	0.5
Resazurin	0.002
Agar-agar**	20.0

^{**}Agar-agar is equivalent to other different terms of agar.

Preparation

Suspend 41.5 g in 1 liter of demineralized water and autoclave (15 minutes at 121 °C).

Cool to about 48 °C and aseptically add, just before use, 5 mL/liter medium freshly prepared ferric (III) ammonium citrate solution (1 g in 5 mL demineralized water, heat sterilized: 15 minutes at 121 °C) and 1.0 mL/liter sodium sulfite solution (1.06657; 2.5 g in 10 mL demineralized water, filter sterilized).

The complete medium is yellowish to reddish-brown.

pH: 7.6 ± 0.2 at 25 °C.

Experimental Procedure and Evaluation

1 mL sample per plate, pour-plate method. After solidification, the plates are overlaid with sterile DCA.

Incubation: At 30 °C for 3 days anaerobically (for example, with Anaerocult[™], Anaerocult[™] A mini).

Reading of results and interpretation: Discrete black colonies of 1-5 mm in diameter are presumptive sulfitereducing clostridia.

Note: To facilitate spore germination, heat treatment of the spores/sample of 10 minutes at 30 °C before inoculation of the agar is recommended.

Storage

Store at 15 °C to 25 °C, dry and tightly closed. Do not use clumped or discolored medium. Protect from UV light (including sunlight). For *in vitro* use only.

The prepared medium is to be used immediately. Do not store.

The base medium can be stored for at least 2 weeks at 4 °C.



Quality Control

Control strains	Growth 48 hours at 30°C, with Anaerocult™ A	Blackening	Recovery
Clostridium perfringens ATCC 10543	Good to Very good	+	≥ 70 %
Clostridium sporogenes ATCC 19404 (WDCM 00008)	Good to Very good	+	≥ 70 %
Clostridium bifermentans ATCC 19299	Good to Very good	+	≥ 70 %
Clostridium perfringens ATCC 13124 (WDCM 00007)	Good to Very good	+	≥ 70 %
Bacillus licheniformis ATCC 14580 (WDCM 00068)	Poor to Fair	-	

Please refer to the actual batch-related Certificate of Analysis.

A recovery rate of 50 % is equivalent to a productivity value of 0.5.

Literature

WEENK, G., FITZMAURICE, E., MOSSEL, D.A.A.: Selective enumeration of spores of *Clostridium* species in dried foods. - J. Appl. Bact., 70; 135-143 (1991).

Ordering Information

Product	Ordering No.	Pack size
Differential Clostridial Agar (DCA) acc. to WEENK	1102590500	500 g
Ammonium iron (III) citrate	F5879-100G	100 g
Sodium sulfite	1066570500	500 g
Anaerocult™ A	1323810001	1 x 10
Anaerocult™ A mini	1323690001	1 x 25

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at <u>SigmaAldrich.com/techservice</u>.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

+Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

Lit No. MS_DS13809EN

MilliporeSigma, Millipore, Anaerocult and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources. © 2024 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

