

## Product Information

**Anti-TNFRSF8**  
antibody produced in chicken,  
affinity isolated antibody

Catalog Number **GW21411A**  
Formerly listed as GenWay Catalog Number 15-288-21411A,  
Tumor necrosis factor receptor superfamily member 8  
Antibody.

Storage Temperature Store at -20 °C

Synonyms: Tumor necrosis factor receptor superfamily,  
member 8 isoform 1, CD30L receptor; Lymphocyte  
activation antigen CD30; KI-1 antigen

### Product Description

Receptor for TNFSF8/CD30L. May play a role in the  
regulation of cellular growth and transformation of  
activated lymphoblasts. Regulates gene expression through  
activation of NF-kappa-B.

NCBI Accession number: NP\_001234.1

Swiss Prot Accession number: P28908

Gene Information: Human .. TNFRSF8 (943)

Immunogen: Recombinant protein Tumor necrosis factor  
receptor superfamily, member 8 isoform 1

Immunogen Sequence: GI # 4507589, sequence 171 - 226

The product is a clear, colorless solution in phosphate  
buffered saline, pH 7.2, containing 0.02% sodium azide.

Species Reactivity: Human

Tested Applications: WB

Recommended Dilutions: Recommended starting dilution  
for Western blot analysis is 1:500, for tissue or cell staining  
1:200.

Note: Optimal concentrations and conditions for each  
application should be determined by the user.

### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or  
other uses. Due to the sodium azide content a material  
safety data sheet (MSDS) for this product has been sent to  
the attention of the safety officer of your institution. Please  
consult the Material Safety Data Sheet for information  
regarding hazards and safe handling practices.

### Storage/Stability

For continuous use, store at 2-8 °C for up to one week. For  
extended storage, store in -20 °C freezer in working  
aliquots. Repeated freezing and thawing, or storage in  
"frostfree" freezers, is not recommended. If slight turbidity  
occurs upon prolonged storage, clarify the solution by  
centrifugation before use. Working dilution samples should  
be discarded if not used within 12 hours.

LPG,MAM12/09-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications.  
Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please  
see reverse side of the invoice or packing slip.