

Product Information

Anti-HOXA13 (N-terminal)

produced in rabbit, affinity isolated antibody

Product Number **H9540**

Product Description

Anti-HOXA13 (N-terminal) is produced in rabbit using as immunogen a synthetic peptide corresponding to a sequence at the N-terminal of human HOXA13 (GeneID: 3209) conjugated to KLH. The corresponding sequence is identical in mouse and rat. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-HOXA13 recognizes human HOXA13 (also known as HOX1, HOX1J). The antibody may be used in several immunochemical techniques including immunoblotting (~40 kDa). Detection of the HOXA13 band by immunoblotting is specifically inhibited with the immunizing peptide.

Hox genes are evolutionarily conserved transcription factors, which act to control important development pathways involved in morphogenesis of the embryo. In vertebrates, there are 39 *HOX* genes that are organized into four clusters (*HOXA–HOXD*) located on different chromosomes (7p15, 17q21.2, 12q13, and 2q31). Each cluster contains 9-11-member genes encoding relatively small gene products. The proteins contain a highly conserved 60-amino-acid region (the homeobox) with DNA-binding activity that contributes to their activity as transcription factors.¹ One of the major functions of *Hox* genes seems to be the formation of the body plan during embryonic development.² In addition to roles in normal development, altered homeobox gene function or expression is implicated in the development of cancers, such as leukemias or in neoplasms of the breast, prostate, kidney, colon, skin and brain.^{3,4} Mutations in the HOXA13 gene are associated with hand-foot-uterus (HFU) syndrome, also known as hand-foot-genital (HFG) syndrome.^{5,6}

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~1.0 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2–8 °C for up to one month. For extended storage, freeze in working aliquots –20 °C. Repeated freezing and thawing, or storage in “frost-free” freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working antibody concentration of 1-2 µg/mL is recommended using lysates of HEK-293T cells over expressing human HOXA13.

Note: In order to obtain best results in various techniques and preparations, it is recommended to determine optimal working dilutions by titration.

References

1. Lemons, D., and McGinnis, W., *Science*, **313**, 1918–1922 (2006).
2. Akam, M., *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, **349**, 313–319 (1995).
3. Stuart, E.T. et al., *Adv. Genet.*, **33**, 255–274 (1995).
4. Cillo, C. et al., *Exp. Cell Res.*, **248**, 1–9 (1999).
5. Stern, A. et al., *Pediatr.*, **77**, 109-116 (1970).
6. Goodman, F. et al., *Am. J. Hum. Genet.*, **67**, 197-202 (2000).

VS,SG,KAA,TD,PHC,MAM 03/19-1