

Product Information

Anti-YB-1 (C-Terminal)

produced in rabbit, affinity isolated antibody

Catalog Number **Y0396**

Product Description

Anti-YB-1 (C-Terminal) is produced in rabbit using as immunogen a synthetic peptide corresponding to amino acids 307-324 of human YB-1 (GeneID: 4904), conjugated to KLH. The corresponding sequence is identical in rat and mouse YB-1 proteins. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-YB-1 (C-Terminal) specifically recognizes YB-1 in human, mouse, and rat. The antibody may be used in several immunochemical techniques including immunoblotting (~50 kDa), immunoprecipitation, and immunofluorescence. Detection of the YB-1 band by immunoblotting is specifically inhibited by the immunizing peptide.

YB-1 (also known as YB1, BP-8, CSDB, DBPB, CSDA2, EFI-A, and NSEP1) is a member of the evolutionarily conserved Cold-Sock Domain (CSD) containing proteins. These are also known as Y-box transcription factors, as they were the first shown to bind the Y-box promoter element to regulate gene expression.¹ These proteins are enriched in the cytoplasm where they are major components of messenger RNP complexes and participate in different steps of mRNA biogenesis. In response to different physiological and environmental stimulation, CDS proteins shuttle between the cytoplasmic and nuclear compartments.^{2,3} Three CDS proteins were identified in mice: YB-1 (encoded by *Ybx1*), YBOX2, known also as DBPC (encoded by *YBX2*) and DBPA (encoded by *Csda*).⁴⁻⁶ Each of these proteins are composed of three domains: the N-terminal which is thought to function as a trans-activation regulatory domain, a highly conserved nucleic acid binding domain (the CSD), and C-terminal domain, thought to be involved in protein-protein interactions.^{2,4}

YB-1 has been shown to play important roles in a wide variety of cellular functions such as transcriptional and translation regulation, DNA repair, drug resistance, and stress responses.⁷⁻⁹ YB-1 is ubiquitously expressed in adult tissues and throughout embryogenesis.

YB-1-deficient embryos die during late embryonic development and exhibit a runting phenotype.⁶

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~1.0 mg/mL

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working antibody concentration of 0.5-1 µg/mL is recommended using PC12 cell lysates.

Immunoprecipitation: a working antibody concentration of 5-10 µg is recommended using HEK-293T cell lysates.

Indirect Immunofluorescence: a working concentration of 2-5 µg/mL is recommended using paraformaldehyde fixed NIH-3T3 cells.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

References

1. Matsumoto, K., et al., *Trends Cell Biol.*, **8**, 318-323 (1998).
2. Kohno, K., et al., *BioEssays*, **25**, 691-698 (2003).

3. Wilkinson, M.F., et al., *BioEssays*, **23**, 775-787 (2001).
4. Davies, H.F., et al., *Dev. Biol.*, **221**, 87-100 (2000).
5. Gu, W.S., et al., *Biol. Reprod.*, **59**, 1266-1274 (1998).
6. Lu, Z.H., et al., *Mol. Cell. Biol.*, **25**, 4625-4637 (2005).
7. Sommerville, J., et al., *BioEssays*, **21**, 319-325 (1999).
8. Evdokimova, V., et al., *J. Biol. Chem.*, **273**, 3574-3581-3832 (1998).
9. Evdokimova, V., et al., *EMBO J.*, **20**, 5491-5502 (2001).

SG,KAA,PHC 06/08-1