

## Product Information

### Anti-G beta ( $G_{\beta}$ )

Developed in Rabbit, Whole Antiserum

Product Number **G4790**

### Product Description

Anti-G beta ( $G_{\beta}$ ) is developed in rabbit using a synthetic peptide (KTREGNVRVSREL) corresponding to a common peptide on G beta ( $G_{\beta}$ ) as immunogen.

Anti-G beta ( $G_{\beta}$ ) is reactive with  $G_{\beta}$  (35-36 kDa) in mammalian tissues by immunoblotting.

Identified by the  $\alpha$  subunit, heterotrimeric G proteins are composed of  $\alpha\beta\gamma$  subunits. They transduce, amplify and diversify the signal generated by the occupancy of a receptor by its hormone or agonist into regulation of one or more effector systems. Receptors activate G proteins by increasing the affinity of the G(GDP) complex for  $Mg^{2+}$ . Bound  $Mg^{2+}$  causes GDP to dissociate and allow GTP to bind and make G(GTP). This is followed by a subunit dissociation reaction that results in active  $\alpha$ (GTP) plus a  $\beta\gamma$  complexed to agonist-occupied receptor. The latter can dissociate further to give free  $\beta\gamma$ . Both the  $\alpha$ (GTP) and  $\beta\gamma$  dimer modulate effector function.

Both  $\alpha$ (GTP) and  $\beta\gamma$  dimers are signaling molecules and modulate positively or negatively a variety of effector functions. The  $\alpha$  subunits remain active until they hydrolyze GTP to GDP.  $\alpha$ (GDP) has high affinity for  $\beta\gamma$  and reconstitutes into trimeric G(GDP) which is ready for another round of nucleotide exchange, activation by GTP and effector modulation through subunit dissociation. Alternatively,  $\alpha$ (GDP) may recombine with  $\beta\gamma$  still associated with the agonist-receptor complex to give a quaternary agonist-receptor-G(GDP) complex ready to be activated. A single ligand occupied receptor is able to activate several G protein molecules during the lifetime of a single  $\alpha$ (GTP) complex. The signal imparted by the binding of a single agonist to its receptor is thus transduced and amplified leading to generation of several active  $\alpha$ (GTP) and  $\beta\gamma$  molecules during the lifetime of the first  $\alpha$ (GTP).

The diversification of the receptor signal is due to: 1) the ability of a single receptor to affect a group of G proteins, such as the  $G_i$ / $G_o$  or the  $G_q$ / $G_{11}$ -class of G proteins; 2)  $\alpha$  and  $\beta\gamma$  subunits having different effects in different cells due to expression of different effectors, and 3) G proteins and their effectors being spatially segregated in a given cell. Thus while in one part of a cell, signaling can occur through  $G_i$  coupled to  $K^+$  channel and adenylyl cyclase (AC), in another part of the cell, responses can be mediated through  $G_o$  coupled to a  $Ca^{2+}$  channel.

$\alpha$  Subunits are encoded in 15 genes and several transcripts are alternatively spliced ( $5\alpha_s$ ,  $2\alpha_{i2}$ ,  $2\alpha_o$  forms). Receptors may discriminate between splice variants; whether splice variants are functionally different in regulating effectors is not known. All  $\alpha$  subunits appear to be palmitoylated near the N-terminus. Palmitate turns over and may affect regulation of GTPase activity by GTPase activating proteins such as regulators of G protein signaling (RGS).

$\beta\gamma$  dimers are heterogeneous and are encoded by 5  $\beta$  and 11  $\gamma$  genes. Some dimers do not form (e.g.,  $\beta_1\gamma_3$ ), and most show tissue selective expression. Structurally,  $\beta$  subunits (four of 340 amino acids and one  $\beta_5$ , of 353 amino acids), are 7-blade propellers, each blade formed of a WD40 motif.  $\gamma$  subunits vary from 68 to 75 amino acids and constitute the most heterogeneous of the three subunit families. All  $\gamma$  subunits are polyiso-prenylated at their C-termini. No functional differences have been found among  $\beta\gamma$  dimers tested with the exception of  $\beta\gamma$  with  $\gamma_1$ , also  $\gamma_T$ , which is less effective probably due to C15-farnesylation instead of C20-geranylgeranylation.  $\beta$  and  $\gamma$  subunits do not exist by themselves, but rather form dimers, or trimers with  $\alpha$  subunits.

### Reagents

Anti-G beta ( $G_{\beta}$ ) is supplied as whole antiserum containing 0.1% sodium azide as a preservative.

### Precautions and Disclaimer

Due to the sodium azide content, a Material Safety Data Sheet (MSDS) for this product has been sent to the attention of the Safety Officer of your institution. Consult the MSDS for information regarding hazardous and safe handling practices.

### Storage/Stability

For continuous use, store at 2-8°C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing is not recommended. Storage in "frost-free" freezers is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

### Product Profile

The recommended dilution is 1:100 to 1:200 for immunoblotting using plasma membrane fractions and chemiluminescence detection.

Note: In order to obtain best results and assay sensitivities of different techniques and preparations, we recommend determining optimal working dilution by titration test.

### References

1. Caro, J.F., et al., J. Cell Biochem., **54**, 309 (1994).
2. De Almeida, J.B., et al., J. Cell Sci., **107**, 507, (1994).
3. Luthin, D.R., et al., J. Biol. Chem., **268**, 5990 (1993).
4. Lynch, et al., J. Clin. Invest., **83**, 2050-2062 (1989).
5. Gettys, et al., J. Biol Chem., **266**, 15949-15955 (1991).
6. Strathmann, M. and Simon, M. I., Proc. Natl. Acad. Sci. USA, **87**, 9113-9117 (1990).
7. Wilkie, T. M. and Yokoyama, K. S. Soc. Gen. Physiol. Ser., **49**, 249-270 (1994).
8. Spiegel, A. M. Mol. Cell Endocrinol., **49**, 1-16 (1987).
9. Roos, W. et al., Biochem. Biophys. Acta, **1448**, 390 (1999).
10. The RBI Handbook of Receptor Classification and Signal Transduction, Watling, K.J., ed., Sigma-Aldrich, p140 (1998).

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.