

Product Information

Anti-Auxin Antibody, Mouse Monoclonal

Clone 1E11-C11, purified from hybridoma cell culture

A0855

Product Description

Anti-Auxin antibody, Mouse monoclonal is derived from the hybridoma 1E11-C11 produced by the fusion of mouse myeloma cells (P3-X63/Ag 8.653) cells and splenocytes from CB6F1/J mice immunized with indoleacetic acid (IAA) carboxyl linked to ovalbumin.¹ The isotype is determined by a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents, Cat. No. ISO2.

Anti-Auxin antibody, Mouse monoclonal recognizes indoleacetic acid but not free unmethylated IAA.¹ The antibody may be used in ELISA,^{1, 3} immunoblotting,¹ dot-blot,¹ and immunohistochemistry.^{2, 4}

Phytohormones play a critical role in plant growth and development. The hormone auxin acts on cells by altering the turgor, elongation, division, and differentiation. Auxin induces the expression of specific genes necessary to regulate growth processes. In most plants, indoleacetic acid is recognized as the key auxin. IAA is synthesized from tryptophan (Trp) using Trp-dependent pathways and from an indolic Trp precursor via Trp-independent pathways.¹⁻²

IAA may be obtained in plants also by β -oxidation of indole-3-butyric acid (IBA) or by hydrolyzing IAA conjugates (IAA may be found linked to amino acids, sugars, or peptides). Conjugation and oxidation of IAA by plants serves to inactivate the hormone. IAA is transported all through the plant (shoot to root) and this transport is important for normal development. Localized transport is needed for tropic responses. The cell signaling of Auxin is mediated through the SCFTIR1 E3 ubiquitin ligase complex. This complex is responsible for the degradation of the transcription repressor Aux/IAA and thereby alters gene expression. The Auxin signal is negatively controlled by auxin-induced genes that encode negatively acting products.¹⁻⁵

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Antibody Concentration: ~2 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free to freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Indirect ELISA: a working antibody concentration of 1-2 µg/mL is recommended using IAA conjugated to BSA as substrate.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.

References

1. Leverone, L.A., et al., Plant Physiol., 96, 1076-1078 (1991).
2. Avsian-Kretchmer, O., et al., Plant Physiol., 130, 199-209 (2002).
3. Pence, V.C., and Caruso, J.L., In: Plant Hormones and their role in plant growth and development, Ed. Davies, P.J., Martinus Nijhoff Publishers, Boston, pp 240-256 (1987).
4. Jiang, K., et al., Development, 130, 1429-1438 (2003).
5. Woodward, A.W., and Bartel, B., Annals Botany, 95, 707-735 (2005).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.
A0855pis Rev 07/22

The Merck logo, consisting of the word "MERCK" in a bold, red, sans-serif font.