# Sigma-Aldrich.

**Technical Bulletin** 

# Anti-ARP3 Antibody, Mouse Monoclonal

Clone FMS338, purified from hybridoma cell culture

#### A5979

## **Product Description**

Anti-ARP3 Monoclonal (mouse IgG2b isotype) is derived from the hybridoma FMS338 produced by the fusion of mouse myeloma cells (NS1 cells) and splenocytes from BALB/c mice immunized with recombinant human ARP3. The isotype is determined by a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents, Cat. No. ISO2.

Anti-ARP3 Monoclonal recognizes human, canine, rat, and mouse ARP3 by ELISA, immunoblotting (40-45 kDa) and immunocytochemistry.<sup>1</sup>

Cell migration processes are mediated by changes in the polymerization state of actin filaments. The signal for these processes begins at the level of the cell surface receptors and is transferred via mediators such as Rac and Cdc42 through Wiskott-Aldrich Syndrome protein (WASP) down to the ARP2/3 complex of proteins. Activation of the actin nucleating activity is mediated by the interaction of domains at the C-terminal part of different WASP family members of proteins with actin-monomers and the ARP2/3 complex. It has been found that ARP2/3 complex and its activator Scar2 are involved in Golgi polarization in NIH 3T3 cells.<sup>2, 3</sup> However in primary astrocytes, Golgi polarization involves neither actin cytoskeleton or Arp2/3 complex nor any WASP-family of proteins.<sup>3</sup> In a different model, it was found that the ARP2/3 complex was necessary for neutrophil chemotaxis and phagocytosis.<sup>4</sup> By genetic and loss of function studies, it was demonstrated that Scar2 is the major regulator of the ARP2/3 complex rather than the WASP family of proteins, the latter having more restricted roles in specific cellular events.5

## Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Antibody concentration: ~ 2 mg/mL

## Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

## Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For prolonged storage, freeze in working aliquots. Repeated freezing and thawing, or storage in frost-free freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

## **Product Profile**

Immunoblotting: A working antibody concentration of 0.25-0.5  $\mu$ g/mL is recommended using total cell extract of mouse myoblast C2 cells.

**Note**: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.



### References

- Baldassarre, M., et al., Eur. J. Cell. Biol., 85, 1217-1231 (2006).
- Magdalena, J., et al., Mol. Biol. Cell, 14, 670-684 (2003).
- Etienne-Manneville, S., and Hall, A., Cell, 106, 489-498 (2001).
- Singh, S., et al., J. Biol. Chem., 278, 36410-36417 (2003).
- Vartiainen, M.K., et al., Curr. Opin. Cell Biol., 16, 174-181 (2004).

#### Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

#### **Technical Assistance**

Visit the tech service page at <u>SigmaAldrich.com/techservice</u>.

#### Standard Warranty

The applicable warranty for the products listed in this publication may be found at <u>SigmaAldrich.com/terms</u>.

#### Contact Information

For the location of the office nearest you, go to <u>SigmaAldrich.com/offices</u>.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

@ 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved. A5979dat Rev07/22

