



Product Information

Ampholine

Product Number **A 5549**
Storage Temperature 2-8 °C

Product Description
CAS Number: 37348-94-0

This product is a sterile filled, 40.0% (w/v) solution. It has been tested to be suitable for use as an isoelectric focusing reagent over the pH range 4.0 - 6.0. The recommend concentration for use is 2%.

Ampholytes are low molecular weight synthetic compounds which are used to generate a pH gradient under application of an electric field. At their respective isoelectric points (pI), the compounds have high buffering capacity. Often referred to as carrier ampholytes, these compounds are commonly used in a defined pH range to give selective resolution to mixtures of amphoteric materials such as proteins. Ampholytes may be used as free reagents or in isoelectric focusing (IEF) strips, whereby the carrier ampholytes are immobilized in a gel matrix.¹

Application of voltage across the ampholyte pH gradient causes the amphoteric molecules of interest to migrate to the gradient positions where their net charges are zero. Materials with net positive charge move toward the cathode, and materials with net negative charge migrate toward the anode. The materials become less charged as they traverse the pH gradient until they reach their respective pI values. Substances which diffuse away from their pI immediately acquire charge and return to their isoelectric position. This focuses the materials at their pI values, separated with very small charge differences. The electric field strength determines the degree of resolution, and thus IEF is performed at high voltages (> 1,000 V). When the materials have reached their final isoelectric positions, the system has very little ionic movement, resulting in a low final current (< 1 mA).

Ampholine® IEF carrier ampholytes are mixtures of 600-700 different homologues within a defined isoelectric point range. Ampholine products are prepared by the reaction of aliphatic oligoamines with acrylic acids. Their molecular weights are generally <1 kDa. Because of their hydrophilic nature, they will not bind to proteins.

Several reports on the use of carrier ampholytes in protein electrophoresis have been published.^{2,3,4,5}

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

References

1. Righetti, P. G., et al., Isoelectric focusing in immobilized pH gradients. *Methods Enzymol.*, **270**, 235-255 (1996).
2. Gorg, A., et al., Recent developments in two-dimensional gel electrophoresis with immobilized pH gradients: wide pH gradients up to pH 12, longer separation distances and simplified procedures. *Electrophoresis*, **20(4-5)**, 712-717 (1999).
3. Gorg, A., et al., The current state of two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis*, **21(6)**, 1037-1053 (2000).
4. Zuo, X., and Speicher, D. W., Quantitative evaluation of protein recoveries in two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis*, **21(14)**, 3035-3047 (2000).
5. Chartogne, A., et al., Capillary electrophoretic separations of proteins using carrier ampholytes. *J. Chromatogr. A*, **959(1-2)**, 289-298 (2002).

Ampholine is a registered trademark of Amersham Biosciences.

GCY/RXR 6/03

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.