

Technical Data Sheet

Sabouraud-2 % Dextrose Broth

acc. harm. EP/USP/JP

Ordering number: 1.08339.0500

Sabouraud Dextrose Bouillon is designed for the cultivation of dermatophytes, yeasts and molds. It can also be used for sterility testing and the determination of the fungi-static activity of pharmaceutical and cosmeceutical products.

This medium complies with the recommendations of the harmonized methods of EP, USP, JP for Microbial Examination of Non-sterile Products: Microbial Enumeration Test and Tests for Specified Microorganisms.

Mode of Action

Sabouraud-2 % Dextrose Broth is used for the cultivation and isolation of yeasts and molds as well as the absence test for *Candida albicans*. The high concentration of dextrose in addition with the low pH promotes the growth, the formation of spores (*Conidia* and *Sporangia*) as well as the formation of pigments of yeasts and molds. On the other side, growth of bacteria is inhibited.

Typical Composition

Peptone from Casein	5 g/l
Peptone from Meat	5 g/l
D(+)-Glucose (= Dextrose)	20 g/l

Preparation

Suspend 30 g/l. If necessary dispense into smaller vessels. Autoclave (15 min at 121 °C).

The appearance of the broth is clear and yellowish-brown.

The pH value at 25 °C is in the range of 5.4-5.8.

Experimental Procedure and Evaluation

Depend on the purpose for which the culture medium is used.

Incubation: up to 7 days at 28 °C aerobically.

Storage

The product can be used for sampling until the expiry date if stored upright, protected from light and properly sealed at +15 °C to +25 °C.

After first opening of the bottle the content can be used up to the expiry date when stored dry and tightly closed at +15 °C to +25 °C.

Disposal

Please mind the respective regulations for the disposal of used culture medium (e.g. autoclave for 20 min at 121 °C, disinfect, incinerate etc.).

Quality Control

Control Strains	ATCC #	Incubation	Expected Results
<i>Trichophyton mentagrophytes</i>	18748	7 days at 28 °C	Growth good to very good
<i>Trichophyton rubrum</i>	28188	7 days at 28 °C	Growth fair to good
<i>Microsporum gallinae</i>	12108	7 days at 28 °C	Growth fair to good
<i>Trichophyton ajelloi</i>	28454	7 days at 28 °C	Growth fair to good
<i>Aspergillus brasiliensis</i> (formerly <i>A. niger</i>)	16404	7 days at 28 °C	Growth good to very good
<i>Penicillium commune</i>	10428	7 days at 28 °C	Growth good to very good
<i>Candida albicans</i>	10231	≤ 3 days at 30-35 °C	Growth fair to very good

Please refer to the actual batch related Certificate of Analysis.

Literature

European Directorate for the Quality of Medicines and Healthcare. (2014): The European Pharmacopoeia. 8th Ed. Chapter 2.6.12 Microbiological examination of non-sterile products: Microbial enumeration tests and Chapter 2.6.13 Microbiological examination of non-sterile products: Test for specified products. Strasbourg, France.

Japanese Ministry of Health, Labour and Welfare. (2011): The Japanese Pharmacopoeia. 16th Ed. Chapter 4.05 Microbial Limit Test I. Microbiological examination of non-sterile products: Total viable aerobic count and II. Microbiological examination of non-sterile products: Test for specified products. Japanese Ministry of Health, Labour and Welfare. Tokyo, Japan.

Mac Faddin, J.J. (1985): Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria. Vol. I. Williams & Wilkins, Baltimore, London, p. 687-691.

Sabouraud R. (1892). Ann. Dermatol. Syphil. **3**: 1061.

United States Pharmacopeial Convention. (2014): The United States Pharmacopeia 38/National Formulation 33, Supp. 2. Chapter <61> Microbiological examination of non-sterile products: Microbial enumeration tests and Chapter <62> Microbiological examination of non-sterile products: Test for specified products. Rockville, Md., USA.



We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

EMD Millipore and the M logo are registered trademarks of Merck KGaA, Darmstadt, Germany. Lit. No. TN1364ENUS

Ordering Information

Product	Cat. No.	Pack size
Sabouraud-2 % Dextrose Broth	1.08339.0500	500 g
Sabouraud-4 % Dextrose Agar	1.05438.0500	500 g
Sabouraud-4 % Dextrose Agar	1.05438.5000	5 kg

EMD Millipore Corporation
290 Concord Road
Billerica, MA 01821, U.S.A.
mibio@emdmillipore.com
www.emdmillipore.com/biomonitoring

Find contact information for your country at:
www.emdmillipore.com/offices
For Technical Service, please visit:
www.emdmillipore.com/techservice



We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

EMD Millipore and the M logo are registered trademarks of Merck KGaA, Darmstadt, Germany. Lit. No. TN1364ENUS