

# Product Information

## AKT2, active, His-tagged, human PRECISIO® Kinase recombinant, expressed in Sf9 cells

Catalog Number **A0858**

Lot Number 011M0787

Storage Temperature –70 °C

Synonyms: PRKBB, PKBBETA, RAC-BETA

### Product Description

AKT2 or Protein Kinase B  $\beta$  (PKB $\beta$ ) is a serine/threonine kinase that is a member of the AKT family. AKT2 like the other AKT members is activated in cells in response to diverse stimuli such as hormones, growth factors, and extracellular matrix components, and is involved in glucose metabolism, transcription, survival, cell proliferation, angiogenesis, and cell motility.<sup>1</sup> The PI3K generates phosphatidylinositol-3,4,5-trisphosphate (PIP<sub>3</sub>), a lipid second messenger essential for the translocation of AKT2 to the plasma membrane, where it is phosphorylated and activated by phosphoinositide-dependent kinase-1 (PDK-1).<sup>2</sup>

This recombinant product was expressed by baculovirus in Sf9 insect cells using an N-terminal His-tag. The gene accession number is NM 001626. It is supplied in 50 mM sodium phosphate, pH 7.0, 300 mM NaCl, 150 mM imidazole, 0.1 mM PMSF, 2 mM DTT, and 25% glycerol.

Molecular mass: ~58 kDa

Purity: ≥70% (SDS-PAGE, see Figure 1)

Specific Activity: 38–52 nmole/min/mg (see Figure 2)


### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.


### Storage/Stability

The product ships on dry ice and storage at –70 °C is recommended. After opening, aliquot into smaller quantities and store at –70 °C. Avoid repeated handling and multiple freeze/thaw cycles.

**Figure 1.**  
SDS-PAGE Gel of Lot Number 011M0787:  
≥70% (densitometry)



**Figure 2.**  
Specific Activity of Lot Number 011M0787:  
40 nmole/min/mg



### Procedure

#### Preparation Instructions

Kinase Assay Buffer – 25 mM MOPS, pH 7.2, 12.5 mM glycerol 2-phosphate, 25 mM MgCl<sub>2</sub>, 5 mM EGTA, and 2 mM EDTA. Just prior to use, add DTT to a final concentration of 0.25 mM.

Kinase Dilution Buffer – Dilute the Kinase Assay Buffer 5-fold with a 50 ng/ $\mu$ l BSA solution.

Kinase Solution – Dilute the active AKT2 (0.1  $\mu$ g/ $\mu$ l) with Kinase Dilution Buffer to the desired concentration.

Note: The lot-specific specific activity plot may be used as a guideline (see Figure 2). It is recommended the researcher perform a serial dilution of active AKT2 kinase for optimal results.

10 mM ATP Stock Solution – Dissolve 55 mg of ATP in 10 ml of Kinase Assay Buffer. Store in 200  $\mu$ l aliquots at –20 °C.

$\gamma$ -<sup>32</sup>P-ATP Assay Cocktail (250  $\mu$ M) – Combine 5.75 ml of Kinase Assay Buffer, 150  $\mu$ l of 10 mM ATP Stock Solution, 100  $\mu$ l of  $\gamma$ -<sup>32</sup>P-ATP (1 mCi/100  $\mu$ l). Store in 1 ml aliquots at –20 °C.

Substrate Solution – Dissolve the synthetic peptide substrate (RPRAATF) in water at a final concentration of 1 mg/ml.

1% phosphoric acid solution – Dilute 10 ml of concentrated phosphoric acid to a final volume of 1 L with water.

#### Kinase Assay

This assay involves the use of the <sup>32</sup>P radioisotope. All institutional guidelines regarding the use of radioisotopes should be followed.

1. Thaw the active AKT2, Kinase Assay Buffer, Substrate Solution, and Kinase Dilution Buffer on ice. The  $\gamma$ -<sup>32</sup>P-ATP Assay Cocktail may be thawed at room temperature.
2. In a pre-cooled microcentrifuge tube, add the following solutions to a volume of 20  $\mu$ l:  
10  $\mu$ l of Kinase Solution  
5  $\mu$ l of Substrate Solution  
5  $\mu$ l of cold water (4 °C)
3. Set up a blank control as outlined in step 2, substituting 5  $\mu$ l of cold water (4 °C) for the Substrate Solution.
4. Initiate each reaction with the addition of 5  $\mu$ l of the  $\gamma$ -<sup>32</sup>P-ATP Assay Cocktail, bringing the final reaction volume to 25  $\mu$ l. Incubate the mixture in a water bath at 30 °C for 15 minutes.
5. After the 15 minute incubation, stop the reaction by spotting 20  $\mu$ l of the reaction mixture onto an individually pre-cut strip of phosphocellulose P81 paper.

6. Air dry the pre-cut P81 strip and sequentially wash in the 1% phosphoric acid solution with constant gentle stirring. It is recommended the strips be washed a total of 3 times of ~10 minutes each.
7. Set up a radioactive control to measure the total  $\gamma$ -<sup>32</sup>P-ATP counts introduced into the reaction. Spot 5  $\mu$ l of the  $\gamma$ -<sup>32</sup>P-ATP Assay Cocktail on a pre-cut P81 strip. Dry the sample for 2 minutes and read the counts. Do not wash this sample.
8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.
9. Determine the corrected cpm by subtracting the blank control value (see step 3) from each sample and calculate the kinase specific activity

#### Calculations:

1. Specific Radioactivity (SR) of ATP (cpm/nmole)

$$SR = \frac{\text{cpm of } 5 \mu\text{l of } \gamma\text{-}^{32}\text{P-ATP Assay Cocktail}}{\text{nmole of ATP}}$$

cpm – value from control (step 7)

nmole – 1.25 nmole (5  $\mu$ l of 250  $\mu$ M ATP Assay Cocktail)

2. Specific Kinase Activity (SA) (nmole/min/mg)

$$\text{nmole/min/mg} = \frac{\Delta \text{cpm} \times (25/20)}{SR \times E \times T}$$

SR = specific radioactivity of the ATP (cpm/nmole ATP)

$\Delta$ cpm = cpm of the sample – cpm of the blank (step 3)

25 = total reaction volume

20 = spot volume

T = reaction time (minutes)

E = amount of enzyme (mg)

#### References

1. Coffer, P.J. et al., Protein kinase B (c-Akt): a multifunctional mediator of phosphatidylinositol 3-kinase activation. *Biochem. J.*, **335**, 1-13 (1998).
2. Anderson, K.E. et al., Translocation of PDK-1 to the plasma membrane is important in allowing PDK-1 to activate protein kinase B. *Curr. Biol.*, **8**, 684-691 (1998).

PRECISIO is a registered trademark of Sigma-Aldrich® Biotechnology LP and Sigma-Aldrich Co.

TD,MAM 01/11-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.