

## GOAT ANTI-MOUSE IgM+IgG+IgA(H+L) FITC CONJUGATED, SPECIES ADSORBED: HUMAN POLYCLONAL ANTIBODY

CATALOG NUMBER: AP501F QUANTITY: 1.0 mg

LOT NUMBER: CONCENTRATION: XX mg/mL

**EPITOPE**: Mouse IgM, IgG, IgA **HOST/ISOTYPE**: Goat

BACKGROUND: IgG is the most abundant immunoglobulin compose of two heavy chains and two light

chains. Each molecule has two antigen binding sites. IgG has 4 subclasses: IgG1 (66%), IgG2 (23%), IgG3 (7%) and IgG4 (4%). IgM constitutes about 10% of serum immunoglobulins. IgM (with IgD) is the major immunoglobulin expressed on the surface of B cells. Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric

IgA is localized to mucosa surfaces.

**SPECIFICITY:** The antibody reacts with the heavy and light chains of mouse IgM, IgG1, IgG2a, IgG2b,

IgG3 and IgA as demonstrated by ELISA and flow cytometry. Minimal cross reactivity with

human immunoglobulins.

**APPLICATIONS:** Immunofluorescence: ≤1 μg/10<sup>6</sup> cells

**ELISA** 

Optimal working dilutions must be determined by the end user.

SPECIES REACTIVITY: Reacts with heavy and light chains of mouse IgM, IgG1, IgG2a, IgG2b, IgG3 and IgA.

Absorbed for human sera and purified human paraproteins. Minimal cross reactivity with

human immunoglobulins. Reactivity with other species has not been determined.

**IMMUNOGEN:** Prepared from pooled antisera from goats hyperimmunized with mouse IgM, IgG and IgA

paraproteins.

**PRESENTATION:** Purified by affinity chromatography on pooled mouse IgM+IgG+IgA covalently linked to

agarose. Liquid in 1.0 mL PBS/NaN<sub>3</sub>.

**STORAGE/HANDLING:** Maintain refrigerated at 2°-8°C under sterile conditions for up to twelve months from date of

receipt.

Important Note: During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For

products with volumes of 200 µL or less, we recommend gently tapping the vial on a hard surface or briefly

centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.

FOR RESEARCH USE ONLY; NOT FOR USE IN DIAGNOSTIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION

Unless otherwise stated in our catalog or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

©2002- 2011: Millipore Corporation. All rights reserved. No part of these works may be reproduced in any form without permission in writing.