

Product Information

Anti-HOXD9 (Internal Region)

produced in rabbit, affinity isolated antibody

Product Number **SAB4200030**

Product Description

Anti-HOXD9 (Internal region) is produced in rabbit using as the immunogen a synthetic peptide corresponding to a fragment of human HOXD9 (Gene ID: 3235) conjugated to KLH. The corresponding sequence is identical in mouse and rat. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-HOXD9 (Internal region) recognizes human HOXD9 (also known as HOX-4C, HOX-5.2). The antibody may be used in several immunochemical techniques including immunoblotting (two bands ~40 kDa in some cell lysate preparations), immunoprecipitation, and immunofluorescence. Detection of the HOXD9 band by immunoblotting is specifically inhibited with the immunizing peptide.

Hox genes are evolutionarily conserved transcription factors, which act to control important development pathways involved in morphogenesis of the embryo. In vertebrates, there are 39 *HOX* genes that are organized into four clusters (*HOXA–HOXD*), located on different chromosomes (7p15, 17q21.2, 12q13, and 2q31). Each cluster contains 9–11 member genes encoding relatively small gene products containing a highly conserved 60-amino-acid region (the homeobox), with DNA-binding activity that contributes to their activity as transcription factors.¹

One of the major functions of *HOX* genes seems to be the formation of the body plan during embryonic development.² In addition to roles in normal development, altered homeobox gene function or expression is implicated in the development of cancers, such as leukemias or neoplasms of the breast, prostate, kidney, colon, skin and brain.^{3,4}

HOXD9 was shown to be involved in the early stages of normal joint development, and also in the pathologic process of arthritis.^{5,6}

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Antibody concentration: ~1.0 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Store at –20 °C. For continuous use, the product may be stored at 2–8 °C for up to one month. For extended storage, freeze in working aliquots at –20 °C. Repeated freezing and thawing, or storage in “frost-free” freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working antibody concentration of 1–2 µg/mL is recommended using lysates of HEK-293T cells overexpressing human HOXD9.

Immunoprecipitation: a working antibody amount of 2.5–5 µg is recommended using lysates of HEK-293T cells overexpressing human HOXD9.

Immunofluorescence: a working antibody concentration of 2–5 µg/mL is recommended using paraformaldehyde fixed HEK-293T cells overexpressing human HOXD9.

Note: In order to obtain best results in various techniques and preparations, it is recommended to determine optimal working dilutions by titration.

References

1. Lemons, D., and McGinnis, W., *Science*, **313**, 1918-1922 (2006).
2. Akam, M., *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, **349**, 313-319 (1995).
3. Stuart, E.T. et al., *Adv. Genet.*, **33**, 255-274 (1995).
4. Cillo, C. et al., *Exp. Cell Res.*, **248**, 1-9 (1999).
5. Khoa, N.D. et al., *Arthritis Rheum.*, **42**, 686-696 (1999).
6. Nguyen, N.C. et al., *Int. J. Mol. Med.*, **10**, 41-48 (2002).

VS,SG,TD,KAA,PHC,MAM 05/19-1