

Product Information

Anti-C-Reactive Protein Antibody, Mouse Monoclonal

Clone CRP-8, Produced from Hybridoma cell culture

C1688

Synonym: Anti-CRP

Product Description

Anti-C-Reactive Protein Antibody, Mouse Monoclonal is derived from the CRP-8 hybridoma produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with purified human C-reactive protein.

Monoclonal Anti-C-Reactive Protein recognizes an epitope located on the 24 kDa subunit of denatured and reduced CRP in an immunoblotting technique. It does not cross react with human serum amyloid P component (SAP), human haptoglobin, human α -1-acid glycoprotein and human IgG, nor with CRP from *Limulus*. The product displays its reactivity against CRP (native and denatured-reduced) in ELISA, dot-blot and immunoblotting.

C-Reactive Protein is a major acute phase reactant synthesized primarily in the liver hepatocytes¹. It is a pentraxin (cyclic pentameric protein) compound of five identical nonglycosylated subunits of 206 amino acids each (24 kDa), that are bound noncovalently to form the physiologic CRP molecule (117.5 kDa). CRP was discovered on the basis of its Ca^{++} ion-dependent binding to the C-polysaccharide of pneumococcal cell wall that occurs via the Ca^{++} dependent site for phosphorylcholine (PC), on each of the subunits of CRP. The CRP molecule has a striking homology with Serum Amyloid P Component (SAP), and regions of sequence homology with the nucleoplasmin/B23 family of proteins.² The precise biological function of CRP is not known. CRP mediates activities associated with preimmune nonspecific host resistance. It is opsonic, an initiator of the classical complement cascade and an activator of monocytes/macrophages.³

Cleavage of CRP by neutrophil-derived proteases leads to the production of peptides that have immunomodulating actions. CRP also binds to several nuclear components including chromatin, histones and snRNP, suggesting that it may play a role as a scavenger during cell necrosis. In addition, CRP binds, through the Ca^{++} -dependent PC-binding site, to many other body components including phospholipids, lecithin, sphingomyelin, a variety of monophosphate esters, low density lipoprotein,⁴ fibronectin and to the basement membrane protein laminin. CRP binding to polycations has also been demonstrated, but this binding does not appear to occur through the PC-binding site on CRP. In humans, normal blood levels of CRP are less than 200 ng/mL, but rise several hundredfold during the first 6-24 hours of an acute inflammation or after tissue injury.

Reagent

Supplied as a solution of 0.01 M Phosphate buffered saline pH 7.4 containing 15 mM Sodium azide as a preservative.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage

For continuous use, store at 2-8 °C. For extended storage, the solution may be frozen in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

Product Profile

Indirect ELISA

A minimum titer of 1:40,000-80,000 was determined using freshly prepared human CRP at 10 µg/mL for coating.

Immunoblotting

A minimum titer of 1:200-400 was determined using human CRP positive serum.

Note: In order to obtain best results in different techniques or preparations, it is recommended that each individual user determine their optimal working dilutions by titration assay.

References

1. Gewurz, H., et al., Adv. Int. Med., 27, 345-372 (1982).
2. DuClos, T.W., et al., J. Immunol., 145, 3869-3875 (1990).
3. Tseng, Y., et al., Hybridoma, 7, 185-191 (1988).
4. Nunomura, W. and Hatakeyama, M., Hokkaido J. Med. Sci., 65, 474-480 (1990).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2008 - 2025 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

23193232 Rev 03/25

