



Product Information

MONOCLONAL ANTI-NICOTINIC ACETYLCHOLINE RECEPTOR, $\alpha 1, \alpha 3, \alpha 5$ SUBUNITS, CLONE mAb210

Purified Rat Immunoglobulin

Product Number **N8033**

Product Description

Monoclonal Anti-Nicotinic Acetylcholine Receptor, $\alpha 1, \alpha 3, \alpha 5$ subunits (rat IgG1 isotype) was produced using denatured nicotinic acetylcholine receptors from bovine and mouse as immunogen¹.

Monoclonal Anti-Nicotinic Acetylcholine Receptor, $\alpha 1, \alpha 3, \alpha 5$ subunits recognizes the main immunogenic region (MIR) of the nicotinic acetylcholine receptor. The antibody detects $\alpha 1$ in all species tested, except *Xenopus*, detects $\alpha 5$ in chicken and human and $\alpha 3$ in human. The antibody passively transfers EAMG.¹ The antibody binds weakly to denatured $\alpha 1$ and strongly with native $\alpha 1$.^{2,3} It does not bind to chicken $\alpha 3, \alpha 4, \alpha 7, \alpha 8, \beta 2$.⁴ The antibody may be used for RIA, immunohistochemistry⁵ and immunoblotting.⁶

Nicotinic acetylcholine receptors (AChR) are members of a gene superfamily of ligand-gated ion channels which includes the homologous GABA_A receptors, glycine receptors and 5-HT₃ serotonin receptors,^{7,8} but not the structurally dissimilar ligand-gated ion channels comprising the glutamate⁹ or ATP receptors.^{10,11} It is likely that all receptors in the AChR superfamily are comprised of five homologous subunits oriented around a central ion channel.¹²

AChR's were first characterized in the skeletal muscles and their structural properties were initially characterized in using AChR's from the homologous electric organ tissue of the *Torpedo* rays,^{12,13,14} The functional and structural characterization of neuronal AChR's developed later due to their lower concentrations in more heterogenous tissues.

Most, if not all, subunits that form the AChR's have now been cloned and expressed. Although more is known now about the structure and function of the neuronal AChR's still little is known about the physiological roles of the many subtypes.

AChR's are now being associated with a growing number of diseases. Thus more research is required to determine the physiological function and role of the

AChR subtypes as well as the receptors themselves in the hopes of discovering new treatments for these pathologies.

Reagents

Monoclonal Anti-Nicotinic Acetylcholine Receptor, $\alpha 1, \alpha 3, \alpha 5$ subunits is supplied at a concentration of approximately 5 mg/ml in 10 mM phosphate buffered saline containing 10 mM sodium azide as a preservative.

Precautions and Disclaimer

Due to the sodium azide content a material safety sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazardous and safe handling practices.

Storage/Stability

For continuous use, store at 2-8°C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing is not recommended. Storage in "frost-free" freezers is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

The recommended working dilution is 1:3,000 – 1:30,000 for immunoblotting and immunohistochemistry.

The moles of *Torpedo californica* electric organ muscle-type receptor α -bungarotoxin binding sites bound per liter of antibody stock solution is measured using a liquid phase RIA as described by Lindstrom, J., et al.,¹⁵ and by a solid phase RIA using goat anti-rat IgG bound to microwell plates. Antibody dilutions for these assays were 1:5,000 – 1:1,000,000. Lot specific titers are given in the certificate of analysis.

Note: In order to obtain best results in different techniques and preparations we recommend determining optimal working dilution by titration test.

References

1. Tzartos, S. et al., *J. Neuroimmunol.*, **15**, 185-194 (1987).
2. Das, M., and Lindstrom, J., *Biochem. Biophys. Res. Commun.*, **165**, 865-871 (1989).
3. Saedi, M.S., *FEBS. Lett.*, **267**, 55 (1990).
4. Conroy, W., et al., *Neuron*, **9**, 1-20 (1992).
5. Keyser, K., et al., *Vis. Neurosci.*, **1**, 349 (1988).
6. Wang, F., et al., *J. Biol. Chem.*, **271**, 17656 (1996).
7. Betz, H., *Neuron*, **5**, 383 (1990).
8. Barnard, E., *Trends Biol. Sci.*, **17**, 368 (1992).
9. Seeburg, P., *Trends Neurosci.*, **16**, 359 (1993).
10. Brake, A., et al., *Nature*, **371**, 519 (1994).
11. Valera, S., et al., *Nature*, **371**, 516 (1994).
12. Lindstrom J. "Ion Channels", Vol. 4, Narahashi, T. (ed), Plenum Press, New York (1996).
13. Changeux, J., Fidia Research Foundation, Neuroscience Award Lectures, **4**, 21 (1990).
14. Karlin, A. and Akabas, M., *Neuron*, **15**, 1231 (1995).
15. Lindstrom, J., et al., Production and Assay of Antibodies to Acetylcholine Receptors, in *Meth. Enzymol.*, vol 74, Langone, J.J., and Van Vunakis, H., (eds.) Academic Press, pp 432-460 (1981).

Sold with the permission of the Salk Institute.

sms/lpg 04/00

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.