

## Product Information

**Anti-BAT3**  
antibody produced in chicken,  
affinity isolated antibody

Catalog Number **GW22823**  
Formerly listed as GenWay Catalog Number 15-288-22823,  
Large proline-rich protein BAT3 Antibody.

Storage Temperature Store at -20 °C

Synonyms: HLA-B associated transcript-3 isoform b, HLA-B-associated transcript 3; Protein G3

### Product Description

A cluster of genes, BAT1-BAT5, has been localized in the vicinity of the genes for TNF alpha and TNF beta. These genes are all within the human major histocompatibility complex class III region. The protein encoded by this gene is a nuclear protein. It has been implicated in the control of apoptosis and regulating heat shock protein.

NCBI Accession number: NP\_542434.1  
Swiss Prot Accession number: P46379

Gene Information: Human .. BAT3 (7917)

Immunogen: Recombinant protein HLA-B associated transcript-3 isoform b

Immunogen Sequence: GI # 18375632, sequence 2 - 250

The product is a clear, colorless solution in phosphate buffered saline, pH 7.2, containing 0.02% sodium azide.

Species Reactivity: Human, rat

Tested Applications: WB

Recommended Dilutions: Recommended starting dilution for Western blot analysis is 1:500, for tissue or cell staining 1:200.

Note: Optimal concentrations and conditions for each application should be determined by the user.

### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Due to the sodium azide content a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

### Storage/Stability

For continuous use, store at 2-8 °C for up to one week. For extended storage, store in -20 °C freezer in working aliquots. Repeated freezing and thawing, or storage in "frostfree" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

LPG,MAM12/09-1