

Product Information

Anti-NONO (N-terminal)

produced in rabbit, affinity isolated antibody

Product Number **N8789**

Product Description

Anti-NONO (N-terminal) is produced in rabbit using as the immunogen a synthetic peptide corresponding to a fragment of human NONO (GenelID: 4841) conjugated to KLH. The corresponding sequence is identical in mouse and rat. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-NONO (N-terminal) recognizes human NONO (also known as p54nrb). The antibody may be used in several immunochemical techniques including immunoblotting (~55 kDa), immunoprecipitation, and immunofluorescence. Detection of the NONO band by immunoblotting is specifically inhibited with the immunizing peptide.

NONO is a nuclear protein implicated in numerous processes including transcription, pre-mRNA processing, nuclear retention of edited RNA, and DNA relaxation.¹ Due to NONO's high similarity to the C-terminus of the human splicing factor PSF, and the fact that it was originally purified as a heterodimer with PSF, NONO was suspected to be involved in pre-mRNA splicing. Later it was found that NONO and PSF mediate different functions depending on their intracellular location.¹

NONO is ubiquitously expressed and composed of two tandem RNP-type RNA-recognition motifs and a putative helix-turn-helix domain followed by a highly charged region.² It is enriched in paraspeckles, a nucleoplasmic compartment, and relocates to cap structures at the nucleolar periphery when transcription is inhibited.³ It was shown that NONO is phosphorylated on multiple sites during mitosis and is a target of the peptidylprolyl isomerase, Pin1.⁴

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~1.0 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Store at -20 °C. For continuous use, the product may be stored at 2–8 °C for up to one month. For extended storage, freeze in working aliquots at -20 °C. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working antibody concentration of 1–2 µg/mL is recommended using NIH-3T3 lysate.

Immunoprecipitation: a working antibody amount of 2.5–5 µg is recommended using HeLa cell lysates.

Immunofluorescence: a working antibody concentration of 2–5 µg/mL is recommended using paraformaldehyde fixed HeLa cells.

Note: In order to obtain best results in various techniques and preparations, it is recommended to determine optimal working dilutions by titration.

References

1. Shav-Tal, Y., and Zipori, D., *FEBS Lett.*, **531**, 109–114 (2002).
2. Dong, B. et al., *Nucl. Acids Res.*, **21**, 4085–4092 (1993).
3. Fox, A. et al., *Curr. Biol.*, **12**, 13–25 (2002).
4. Proteau, A. et al., *J. Mol. Biol.*, **346**, 1163–1172 (2005).

VS,SG,TD,KAA,PHC,MAM 04/19-1