

Data Sheet

LX-2 Cas9 Human Hepatic Stellate Cell Line

Immortalized Cell Line

SCC613

Pack Size: $\geq 1 \times 10^6$

Store in Liquid Nitrogen

FOR RESEARCH USE ONLY**Not for use in Diagnostic Procedures. Not for Human or Animal Consumption.**

Background

Hepatic stellate cells perform vital roles in response to injury in the liver and are the body's major storage site for vitamin A. However, activation of hepatic stellate cells into fibrogenic myofibroblast-like cells in response to infection such as hepatitis and chronic conditions such as chronic alcoholism, fatty liver disease, obesity and diabetes causes liver fibrosis. Advanced liver fibrosis results in cirrhosis and liver failure. Given the importance of stellate cells in disease etiology, genetically malleable cellular models are critical for understanding the contribution of these cells to liver pathologies and necessary for development of new therapeutics.

The immortalized LX-2 cell line was established by Xu et al. to provide a stable, homogenous, and unlimited source of human hepatic stellate cells and overcome issues with culture variability. Since its introduction, LX-2 has proven a successful and physiologically relevant model for both clinical and basic research and development, as LX-2 cells retain key features of cytokine signaling, neuronal gene expression, retinoid metabolism, and fibrogenesis. The Cas9 version of the LX-2 cell line has been developed to allow targeted CRISPR/Cas9 modification and knockouts of genes in LX-2 cells. LX-2 Cas9 human hepatic stellate cells express a Cas9 cassette and transfection with CRISPR-cassettes to genes of interest enables direct assessment of genes associated with fibrogenesis and stellate cell disease response. The LX-2 Cas9 hepatic stellate cell line expresses common stellate cell markers such as MMP2 and alpha-smooth muscle actin while exhibiting slightly slower growth than original LX-2 cells in low-serum media.

Products in this document can be purchased at [SigmaAldrich.com](https://www.sigmaaldrich.com) using the catalogue numbers in parenthesis unless otherwise noted.

Source

LX-2 Cas9 human hepatic stellate cells were derived from LX-2 cells transformed with a Cas9 cassette.

Short Tandem Repeat (STR) Profile

D3S1358: 13, 15	D21S11: 28, 31	TH01: 9.3	Penta D: 13, 15
D7S820: 11	D18S51: 12	TPOX: 8, 9	Penta E: 5, 21
vWA: 17	D5S818: 11, 12	CSF1P0: 10, 12	
FGA: 13	D13S317: 11, 13	AMEL: X, Y	
D8S1179: 13	D16S539: 13		

Quality Control Testing

- Each vial contains $\geq 1 \times 10^6$ viable cells.
- Cells are tested negative for infectious diseases by a Human Essential Cell Line Examination and Report (CLEAR) Panel by Charles River Animal Diagnostic Services.
- Cells are verified to be of human origin and negative for inter-species contamination from mouse, rat, Chinese hamster, Golden Syrian hamster, and Non-human Primate (NHP) as assessed by a Contamination CLEAR Panel by Charles River Animal Diagnostic Services.
- Cells are negative for mycoplasma contamination.

Storage and Handling

LX-2 Cas9 human hepatic stellate cells should be stored in liquid nitrogen. The cells can be cultured for at least 10 passages after initial thawing without significantly affecting functionality.

Representative Data





Figure 1: A. Bright-field image of LX-2 Cas9 cells one day after thaw in a T75 flask. B. Cell counting was performed using Scepter™ 3.0 Handheld Automated Cell Counter using 60 μm sensor tips (PHCC360KIT). C. LX-2 Cas9 cells express α-smooth muscle actin (SMA), D. matrix metalloproteinase 2 (MMP2) and E. Vimentin.

Figure 2: Schematic map of LentiCas9-Blast Vector



Protocol

Thawing Cells

1. Do not thaw the cells until the recommended medium is on hand. Cells can grow on normal tissue cultureware surfaces without any additional coating.
Cells are thawed and expanded in LX-2 Cas9 Expansion Medium comprised of DMEM-High Glucose medium (D5796) with 10% FBS (ES-009-B), 2 mM L-Glutamine (TMS-002-C), and 1X penicillin/streptomycin (TMS-AB2-C, optional).
2. Remove the vial of frozen LX-2 Cas9 cells from liquid nitrogen and incubate in a 37 °C water bath. Closely monitor until the cells are completely thawed. Maximum cell viability is dependent on the rapid and complete thawing of frozen cells.
IMPORTANT: Do not vortex cells.
3. As soon as the cells are completely thawed, disinfect the outside of the vial with 70% ethanol. Proceed immediately to the next step.
4. In a laminar flow hood, use a 1 mL or 2 mL pipette to transfer the cells to a sterile 50 mL conical tube. Be careful not to introduce any bubbles during the transfer process.
5. Using a 10 mL pipette, slowly add dropwise 9 mL of LX-2 Cas9 Expansion Medium (Step 1 above) to the 15 mL conical tube.
IMPORTANT: Do not add the entire volume of media all at once to the cells. This may result in decreased cell viability due to osmotic shock.
6. Gently mix the cell suspension by slowly pipetting up and down twice. Be careful not to introduce any bubbles.
IMPORTANT: Do not vortex cells.
7. Centrifuge the tube at 300 x *g* for 2-3 minutes to pellet the cells.
8. Decant as much of the supernatant as possible. Steps 5-8 are necessary to remove residual cryopreservative (DMSO).
9. Resuspend the cells in 15 mL of LX-2 Cas9 Expansion Medium.
10. Transfer the cell mixture to a T75 tissue culture flask.
11. Incubate the cells at 37 °C in a humidified incubator with 5% CO₂.

Subculturing Cells

1. LX-2 Cas9 cells should be passaged at ~80-85% confluency. Do not allow the cells to grow over 85% confluency.
2. Carefully remove the medium from the T75 flask containing the 80% confluent layer of LX-2 Cas9 cells. Rinse the flask with 10 mL 1X PBS. Aspirate after the rinse.
3. Apply 3-5 mL of Accutase® or Trypsin/EDTA solution and incubate in a 37 °C incubator for 3-5 minutes.
4. Inspect the flask and ensure the complete detachment of cells by gently tapping the side of the flask with the palm of your hand.
5. Add 8 mL of LX-2 Cas9 Expansion Medium to the plate.
6. Gently rotate the flask to mix the cell suspension. Transfer the dissociated cells to a 15 mL conical tube.
7. Centrifuge the tube at 300 x *g* for 3-5 minutes to pellet the cells.
8. Discard the supernatant, then loosen the cell pellet by tapping the tip of the tube with a finger.
9. Apply 2-5 mL of LX-2 Cas9 Expansion Medium to the conical tube and resuspend the cells thoroughly.
IMPORTANT: Do not vortex the cells.
10. Count the number of cells using a hemocytometer or a Scepter™ 3.0 Handheld Automated Cell Counter using 40 µm sensor tips.
11. Plate the cells to the desired density. Typical split ratio is 1:5 to 1:6. The medium should be replaced every other day.

Cryopreservation of Cells

LX-2 Cas9 human hepatic stellate cells may be frozen in LX-2 Cas9 Expansion Medium and 10% DMSO using a Nalgene™ slow freeze Mr. Frosty® container.

References

1. *Gut* 2005, 54(1): 142-151.
2. *PloS One* 2013, 8(10): e75692.

Academic Use Agreement

(Subject to local law)

THIS PRODUCT MAY ONLY BE USED BY INDIVIDUALS EMPLOYED BY AN ACADEMIC INSTITUTION AND IS INTENDED SOLELY TO BE USED FOR ACADEMIC RESEARCH, WHICH IS FURTHER DEFINED BELOW. BY OPENING THIS PRODUCT, YOU ("PURCHASER") HEREBY REPRESENT THAT YOU HAVE THE RIGHT AND AUTHORITY TO LEGALLY BIND YOURSELF AND/OR YOUR EMPLOYER INSTITUTION, AS APPLICABLE, AND CONSENT TO BE LEGALLY BOUND BY THE TERMS OF THIS ACADEMIC USE AGREEMENT. IF YOU DO NOT AGREE TO COMPLY WITH THESE TERMS, YOU MAY NOT OPEN OR USE THE PRODUCT AND YOU MUST CALL CUSTOMER SERVICE (1-800-645-5476) TO ARRANGE TO RETURN THE PRODUCT FOR A REFUND.

"Product" means LX-2 Cas9 Human Hepatic Stellate Cell Line (SCC613).

"Academic Research" means any internal in vitro research use by individuals employed by an academic institution. Academic Research specifically excludes the following uses of whatever kind or nature.

- Re-engineering or copying the Product
- Making derivatives, modifications, or functional equivalents of the Product
- Obtaining patents or other intellectual property rights claiming use of the Product
- Using the Product in the development, testing, or manufacture of a Commercial Product
- Using the Product as a component of a Commercial Product
- Reselling or licensing the Product
- Using the Product in clinical or therapeutic applications including producing materials for clinical trials
- Administering the Product to humans
- Using the Product in collaboration with a commercial or non-academic entity

"Commercial Product" means any product intended for: (i) current or future sale; (ii) use in a fee-for-service; or (iii) any diagnostic, clinical, or therapeutic use.

Access to the Product is limited solely to those officers, employees, and students of PURCHASER's academic institution who need access to the Product to perform Academic Research. PURCHASER shall comply with all applicable laws in its use and handling of the Product and shall keep it under reasonably safe and secure conditions to prevent unauthorized use or access.

These use restrictions will remain in effect for as long as PURCHASER possesses the Product.

This Product and its use may also be covered by one or more patents used under license from owners or licensors of CRISPR intellectual property found here: <https://www.sigmaldrich.com/US/en/life-science/legal/product-licenses/crispr-use-license-agreement>.

COMMERCIAL OR NON-ACADEMIC ENTITIES INTERESTED IN PURCHASING OR USING THE PRODUCT MUST CONTACT licensing@emdmillipore.com AND AGREE TO SEPARATE TERMS OF USE PRIOR TO USE OR PURCHASE.

GMO

This product contains genetically modified organisms.

Este producto contiene organismos genéticamente modificados.

Questo prodotto contiene degli organismi geneticamente modificati.

Dieses Produkt enthält genetisch modifizierte Organismen.

Ce produit contient des organismes génétiquement modifiés.

Dit product bevat genetisch gewijzigde organismen.

Tämä tuote sisältää geneettisesti muutettuja organismeja.

Denna produkt innehåller genetiskt ändrade organismer.

Denna produkt innehåller genetiskt ändrade organismer.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany
operates as MilliporeSigma in the U.S. and Canada.

Merck, Scepter and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2024 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

Document Template 20306518 Ver 6.0

23027675 Ver 1.0, Rev 02FEB2024, AB

