



**RABBIT ANTI-BRAIN DERIVED NEUROTROPHIC FACTOR
(IgG Fraction)
POLYCLONAL ANTIBODY**

CATALOG NO:	AB1534SP
LOT NUMBER:	xxx
QUANTITY:	500 µg
SPECIFICITY:	Brain Derived Neurotrophic Factor (BDNF). By dot blot, less than 0.1% cross-reactivity against mouse NGF, recombinant human NT3 or neurotrophin 4.
IMMUNOGEN:	A 10 amino acid peptide corresponding to the amino-terminal of mouse Brain Derived Neurotrophic Factor (BDNF) coupled to ovalbumin.
APPLICATIONS:	Immunohistochemistry: 1-10 µg/mL (see suggested protocol). Immunoblotting: 1-10 µg/mL ELISA (one-site): 1-10 µg/mL Optimal working dilutions must be determined by end user.
SPECIES REACTIVITIES:	Rat and human.
FORMAT:	Protein A purified immunoglobulin
PRESENTATION:	Lyophilized from PBS, no preservatives. Reconstitute with 500 µL of sterile distilled water (1 mg/mL).
STORAGE/HANDLING:	Maintain lyophilized material at -20 to -70°C for up to 12 months after date of receipt. After reconstitution maintain at -20 to -70°C in undiluted aliquots for up to 6 months. Avoid repeated freeze/thaw cycles.

NEUROTROPHIN IMMUNOHISTOCHEMISTRY

The neurotrophins have proved difficult to localize which may be due to masking by, for example, their association with the *trk* receptors or very low concentrations. Where success has been achieved, the conditions required vary greatly for different tissues. A protocol has been included for central nervous system tissue and, while it is similar to commonly used methods, it is important to give strict attention to details such as thorough washing, fixative and detergent concentrations, concentration and quality of the primary antibodies and length of incubations, etc. You may find it possible to use alternate protocols, however we have experienced many failures using variations of the current protocol (and some failures when strict adherence to the procedure is maintained). A suitable procedure to stain nerve terminals is still being developed. Neurotrophin receptors have proved much easier to localize.

It is recommended that you include a few sections of adult rat cerebellum, spinal cord or kidney in each experiment since these are tissues which are the easiest to stain.

Protocol for Immunohistochemistry in the Central Nervous System

At all steps thorough washing is necessary to reduce background.

Fixation

Animals are perfused with 1% sodium nitrite in phosphate buffered saline (PBS) (about 50 mL) followed by 1 liter of Zamboni's fixative (4% formaldehyde, 15% picric acid in 0.1M phosphate buffer). Post fix for no longer than 2 hours.

Tissue Preparation

Tissues are removed and washed briefly with PBS followed by cryoprotection in 30% sucrose in PBS overnight at 4°C. 30 µm cryostat sections are cut and washed with agitation in:

PBS	(1x15min)
50% ethanol	(3x15min)
PBS	(1x15min)

Sections can be stored at 4°C in TBS for several weeks in the presence of 0.2% sodium azide.

Blocking and Primary Antibody Incubations

A 24 well tissue culture plate works well for incubations. Sections are blocked with 20% normal horse serum in PBS for at least 1 hour followed by incubation with primary antibody diluted in 2xPBS, 0.3% Triton X-100 containing 0.02% sodium azide. Incubation can range from 24 hrs to 1 week. Room temperature (RT) is used for 24 hr incubations but 4°C is used for longer incubations.

Neurotrophin antibodies are normally used after affinity purification at a concentration of approximately 0.5-1.0 µg/mL. Prepare sufficient antibody only to cover the sections.

Secondary Antibodies

Primary antibodies are removed and sections are washed in PBS-T (PBS + 0.1% Tween), 3x15min. Biotinylated affinity purified IgG antibody is then applied to the sections for 2 hours at RT followed by another 3x15 min washes in PBS-T.

ABC Reagent: The ABC reagent is prepared 30 minutes prior to use and applied to sections for 2 hours at RT according to the manufactures recommendation.



ABC reagent is then removed and sections are washed with Tris buffered saline (TBS), 3x15 mins, to remove all traces of the ABC.

Development

0.25% nickel sulfate/TBS solution is prepared and to 20 mls of this solution, 10 mg Diaminobezidine (DAB) is added. Immediately prior to use, 7.5 μ L of hydrogen peroxide (30% w/v) is added to this solution. Sections are incubated in this solution for up to 30 minutes, until the bluish color develops. If sections show rapid color change due to high background, the neurotrophin immunoreactivity will be difficult to detect. Washing procedure will then need to be improved.

To end the reaction, remove the DAB solution and wash sections in TBS, 3x15 mins. Transfer sections to glass microscope slides, stretch and arrange using a small paintbrush. Slides are then air dried, dehydrated through graded alcohols, cleared in xylene, and mounted in a xylene based mounting media.

Sections can then be examined using light microscopy.

Protocol for Neurons in the Peripheral Nervous System

See Zhou et al., (1994) *J. Neuroscience Methods* **54**:95-102. This protocol works well for neurons in cranial and spinal ganglia as well as for those in sympathetic ganglia.

Important Note: *During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 μ L or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.*

FOR RESEARCH USE ONLY; NOT FOR USE IN DIAGNOSTIC
PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION

Unless otherwise stated in our catalog or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

©2003 - 2011: Millipore Corporation. All rights reserved. No part of these works may be reproduced in any form without permission in writing.

28820 Single Oak Drive • Temecula, CA 92590
Technical Support: T: 1-800-MILLIPORE (1-800-645-5476) • F: 1-800-437-7502
www.millipore.com