

Product Information

Monoclonal Anti-Human CD20 Clone B9E9

Purified Mouse Immunoglobulin

Product Number **C 8080**

Product Description

Monoclonal Anti-Human CD20 (mouse IgG2a isotype) is derived from the B9E9 hybridoma produced by the fusion of mouse myeloma cell line NS-1 and splenocytes from an immunized mouse. The human lymphoblastoid cell line Daudi was used as immunogen.¹ The isotype is determined using Sigma ImmunoType™ Kit (Sigma ISO-1) and by a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents (Sigma ISO-2).

Monoclonal Anti-Human CD20 recognizes the B cell-specific antigen, human CD20, expressed by pre-B cells, resting, and activated B cells but not by pro-B cells, plasma cells, or myeloma cells. The antibody may be used in flow cytometry, immunohistochemistry (frozen sections), and immunocytochemistry. The epitope recognized by the antibody is sensitive to routine formalin fixation and paraffin embedding procedures. In flow cytometric analysis, the antibody reacts dimly with T cells, thus requiring resetting the boundary between positive and negative cells to include only the brightly stained cells as positive.

CD20 is a 33/35/37 kDa membrane-embedded, unglycosylated phosphoprotein with four membrane-spanning domains as well as serine and threonine rich cytoplasmic N-terminal and C-terminal domains. Only a minor portion of the molecule consisting of one short and one longer loop is exposed on the cell surface.²⁻⁴ CD20 exists on the cell surface as a homo-oligomeric complex. CD20 is expressed on some non T-ALL cells and possibly on follicular dendritic cells but not on monocytes, macrophages, neutrophils, red blood cells, or platelets. In normal lymph nodes and tonsils, CD20 is detected on B cells in germinal centers while lower density expression is manifested on cells in the mantle zone.⁵ CD20 is associated with transmembrane Ca^{2+}

conductance and is involved in the regulation of B cell proliferation and differentiation. While CD20 is not phosphorylated in resting B cells, it is differentially phosphorylated in its cytoplasmic domains by ubiquitous kinases in activated cells or B cell lines. Binding of many monoclonal antibodies to CD20 may cause the generation of transmembrane signals that result in enhanced CD20 and other cellular proteins phosphorylation, induction of oncogene expression, and increased expression of CD18, CD58, and MHC Class II antigens.⁶ Other antibodies may inhibit B cell proliferation and differentiation.

Monoclonal Anti-Human CD20 may be used for:

1. Identification of normal, leukemic, and lymphoma B cells in smears, cytopins, cell suspensions, and frozen sections.
2. Enumeration of B lymphocytes in peripheral blood.
3. B cell isolation by sorting procedures.

Reagent

The product is supplied as purified antibody in a solution containing 0.01 M phosphate buffered saline, pH 7.4, containing 1% bovine serum albumin and 15 mM sodium azide as a preservative.

Precautions and Disclaimer

Due to the sodium azide content, a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazardous and safe handling practices.

Storage/Stability

Store at 2-8 °C. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

By indirect immunofluorescence, a working antibody dilution of 1:10 is recommended using human tonsil frozen sections.

By flow cytometry, 5 μ l of the monoclonal antibody will stain 1×10^6 cells with a fluorescence intensity and percent positive that is similar to that observed with saturating amounts of monoclonal antibody.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.

Procedure for Indirect Immunofluorescent Staining

Reagents and materials needed but not supplied:

- Whole human blood collected by standard clinical blood evacuation tubes with EDTA, ACD-A, or heparin anticoagulant **or**
 - Human cell suspension (e.g. peripheral blood mononuclear cells isolated on HISTOPAQUE[®] (Sigma Stock No. 1077-1)).
- Diluent (freshly prepared): 0.01 M phosphate buffered saline (PBS), pH 7.4, containing 1% BSA.
- Fluorochrome (FITC, PE, or Quantum Red[™]) conjugated anti-mouse secondary antibody diluted to recommended working dilution in diluent (e.g. Product No. F2883 FITC-Sheep Anti-Mouse IgG (whole molecule), F(ab')₂ fragment of affinity isolated antibody). Aggregates in conjugates should be removed by centrifugation immediately prior to use.
- Isotype-matched, non-specific mouse immunoglobulin (negative control, Product No. M5409).
- 12 x 75 mm test tubes.
- Adjustable micropipet.
- Centrifuge.
- Counting chamber.
- 0.2% Trypan blue (Product No. 302643) in 0.01 M phosphate buffered saline, pH 7.4.
- 2% paraformaldehyde in PBS.
- Whole blood lysing solution.
- Flow cytometer.

Assay procedure

- Use 100 μ l of whole blood **or**
 - Adjust cell suspension to 1×10^7 cells/ml in diluent. Cells should be >90% viable as determined by dye exclusion (e.g., trypan blue). For each sample, add 100 μ l or 1×10^6 cells per tube.
- Add 5 μ l of monoclonal antibody to tube(s) containing cells to be stained. Vortex tube gently. Incubate the cells at room temperature for 30 minutes.
Proper controls to be included for each sample are:
 - Autofluorescence control: 5 μ l diluent in place of monoclonal antibody.
 - Negative staining control: 5 μ l isotype-matched non-specific mouse immunoglobulin (Product No. M5409) at the same concentration as test antibody.
- After 30 minutes, add 2 mL of diluent to all tubes.
- Pellet cells by centrifugation at 500 x G for 10 minutes.
- Remove supernatant by careful aspiration.
- Resuspend cells in 2 mL diluent.
- Repeat washing procedure (steps 4-6).
- After the second wash, resuspend the cells in 100 μ l of the fluorochrome conjugated secondary antibody at the recommended concentration. For the autofluorescence control, add 100 μ l of diluent. Incubate at room temperature for 30 minutes. Protect from light at this and all subsequent steps.
- If whole blood is used, use lysing solution after incubation according to manufacturer's instructions, then proceed to Step 8.
 - If a mononuclear cell suspension is used, proceed to Step 8.
- Add 2 mL diluent to all tubes.
- Wash as in steps 4-6 twice.
- After last wash, resuspend cells in 0.5 mL of 2% paraformaldehyde and analyze in a flow cytometer according to manufacturer's instructions.

Quality Control

It is advisable to run the appropriate negative controls. Negative controls establish background fluorescence and non-specific staining of the primary and secondary antibodies. The ideal negative control reagent is a mouse monoclonal or myeloma protein. It should be isotype-matched to the primary antibody, not specific for human cells and of the same concentration as the primary antibody. The degree of autofluorescence or negative control fluorescence will vary with the type of cells under study and the sensitivity of the instrument used. For fluorescence analysis of cells with Fc receptors, the use of isotype matched negative control is mandatory. In some systems it may be necessary to incubate the cells in 10-20% normal serum from the second antibody host species (at step 2 before adding monoclonal antibody) in order to decrease non-specific staining with the conjugated second antibody.

References

1. Leucocyte Typing V, Schlossman, S., et al., (eds.), Oxford University Press, Oxford, pp 511, 513, 515 (1995).
2. Tedder, T., et al., J. Immunol., **141**, 4388 (1988).
3. Tedder, T., et al., Proc. Natl. Acad. Sci., USA, **85**, 208 (1988).
4. Tedder, T., et al., J. Immunol., **142**, 2560 (1989).
5. Ledbetter, J., and Clark, E., Human Immunology, **15**, 30 (1986).
6. Tedder, T., and Engel, P., Immunology Today, **15**, 450 (1994).

KAA 10/05

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.