

Product Information

Anti-Strumpellin (C-terminal)

produced in rabbit, affinity isolated antibody

Catalog Number **SAB4200550**

Product Description

Anti-Strumpellin (C-terminal) is produced in rabbit using as immunogen a synthetic peptide corresponding to the C-terminal region of human Strumpellin (GenelD: 9897), conjugated to KLH. The corresponding sequence is identical in mouse and rat. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-Strumpellin (C-terminal) recognizes human Strumpellin. The antibody may be used in various immunochemical techniques including immunoblotting (~110 kDa) and immunoprecipitation. Detection of the Strumpellin band by immunoblotting is specifically inhibited by the immunizing peptide.

Strumpellin is a component of the WASH multiprotein complex, an actin-regulating complex that is recruited to endosomes by interactions with the retromer complex. In addition to Strumpellin, the WASH complex is composed of WASH1, FAM21, KIAA1033 (SWIP) and CCDC53. Mutations in the gene encoding strumpellin, *KIAA0196*, cause hereditary spastic paraplegia, a progressive neurodegenerative disorder clinically characterized by central motor system deficits leading to spastic paraparesis of the lower limbs. Strumpellin presence was observed in protein aggregate diseases affecting striated muscles and the CNS.¹⁻⁵

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody Concentration: ~ 1.0 mg/mL

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage freeze in working aliquots. Repeated freezing and thawing is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working concentration of 5-10 µg/mL is recommended using whole extracts of human PC-3 cells.

Immunoprecipitation: a working amount of 5-10 µg is recommended using lysates of human DU-145 cells.

Note: In order to obtain best results in different techniques and preparations we recommend determining optimal working concentration by titration test. Exposure to sensitive film is recommended.

References

1. Valdmanis, P.N., et al., *Am. J. Hum. Genet.*, **80**, 152-161 (2007).
2. Jia, D., et al., *Proc. Natl. Acad. Sci. USA*, **107**, 10442-10447 (2010).
3. Harbour, M.E., et al., *J. Cell Sci.*, **123**, 3703-3717 (2010).
4. Clemen, C.S., et al., *Brain*, **133**, 2920-2941 (2010).
5. Freeman, C., et al., *Biochim. Biophys. Acta*, (2012) [Epub ahead of print].

ST,KCP,PHC 01/13-1