

Test Agar pH 6.0 for the Inhibitor Test

For the detection of antimicrobial inhibitors in meat and organ samples together with *Bacillus subtilis* (BGA) spore suspension and *Micrococcus luteus* ATCC 9341 as test organisms.

The nutrient media are suitable both, for the inhibitor test (LEVETZOW, 1971) according to the German Meat Inspection Law as well as for the EEC Four-Plate-Test (BOGAERTS and WOLF, 1980) suggested by the Scientific Veterinary Commission of the European Economic Community. Test agar pH 7.2 with addition of trimethoprim is used particularly for determination of sulfonamide residues.

Mode of Action

The test is carried out according to the agar diffusion procedure. Small slices of the meat sample are placed on the inoculated Test Agar plates and incubated. Antimicrobial inhibitors contained in the samples diffuse into the nutrient media and cause growth-free inhibition zones to develop on the otherwise thickly covered plates. Repeated tests with pH 6.0, pH 8.0 and pH 7.2 are necessary, as penicillin and streptomycin are optimally active at pH 6.0 and 8.0 respectively (PICHNARCIK et al., 1969) and the activity optimum of sulfonamide is found at pH 7.2. Addition of trimethoprim to Test Agar pH 7.2 considerably increases the sensitivity of the test system to sulfonamides (GUDDING, 1976; EBRECHT, 1982).

Typical Composition (g/litre)

Peptone from casein, tryptic 3.45; peptone from meat, tryptic 3.45; sodium chloride 5.1; agar-agar 13.0.

Preparation

Suspend 25 g/litre Test Agar pH 6.0, autoclave (15 min at 121 °C), test the pH and if necessary adjust. Cool to 50-45 °C, mix in 1ml/litre *Bacillus subtilis* (BGA) spore suspension.

After mixing the spore suspension, immediately pour the plates and place in the refrigerator.

The plates are clear and yellowish-brown.

Storage of the Plates

The ready-to-use plates can be sealed with air-tight adhesive tape and, when cooled (+4 to +6 °C) may be kept for up to 2 weeks. Additional packing into plastic bags is recommended. If stored for longer periods, the temperature should not be more than +3 °C; however, freezing of the culture medium must be avoided.

Experimental Procedure and Evaluation

Inhibitor Test

Test Agar pH 6.0 with *Bacillus subtilis* (BGA).

EEC Four-Plate-Test

Test Agar pH 6.0 with *Bacillus subtilis* (BGA).

For details regarding the collection of samples, transportation as well as the execution of the test, see the Fleischbeschaugesetz (Meat Inspection Law) or BOGAERTS and WOLF (1980).

According to these specifications the cylinder-shaped tissue sections (8 mm in diameter and 2 mm thick) are stamped out under conditions, which are as aseptic as possible and laid on the plates; according to BOGAERTS and WOLF two sections are required per plate. As a control, one test disc with 10 IU of penicillin G-sodium is placed on a plate with pH 6.0, one test disc with 10 µg of streptomycin on each of the two plates with pH 8.0 and one test disc with 0.5 µg sulfadimidine on a plate with pH 7.2. The test discs can also be made by the user using filter paper discs, 6 mm in diameter.

Incubation: 18-24 hours at 30 °C (*Bacillus subtilis* BGA).

Manufacturers	Product
Burroughs Wellcome, GB	Trimethoprim
Intern. Chemical Industries, GB	Sulfadimidine
Schleicher & Schüll, No. 2668, FRG	Filter paper discs 6 mm in diameter
American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, USA	<i>Micrococcus luteus</i> ATCC 9341
Beiersdorf AG, Hamburg, FRG	Adhesive tape Tesaflex 166

The inhibition zones between the edge of the tissue section or the test disc and the growth limit of the test organism are measured. Complete inhibition of growth with an inhibitory zone of at least 2mm is regarded as positive result, an inhibitory zone of 1-2mm should be considered as doubtful. This only applies, however, if the controls, prepared at the same time, display zones of inhibition measuring about 6 mm.

For possible methodological improvements see FORSCHNER and SEIDLER (1976).

Test Agar pH 6.0 for the Inhibitor Test

Literature

Arbeitsgruppe des Instituts für Lebensmitteltechnologie und Verpackung der Technischen Universität München: Merkblätter für die Prüfung von Packmitteln, Merkblatt 18 "Prüfung auf antimikrobielle Bestandteile in Packstoffen". - **Verpackgs.-Rdsch.**, **25**; Techn.-wiss. Beilagen; 5-8 (1974).

BAUR, E.: Untersuchungen von Fleisch- und Wurstwaren mit dem Hemmstofftest im Rahmen der tierärztlichen Lebensmittelüberwachung. - **Fleischwirtsch.**, **55**; 843-845 (1975)

BOGAERTS, R., u. WOLF, F.: Eine standardisierte Methode zum Nachweis von Rückständen antibakteriell wirksamer Substanzen in frischem Fleisch. - **Fleischwirtsch.**, **60**; 667-675 (1980).

Deutsches Fleischbeschaugesetz: Ausführungsbestimmungen A über die Untersuchung und gesundheitspolizeiliche Behandlung der Schlachttiere und des Fleisches bei Schlachtungen im Inland; Anlage 4 zu § 20 Abs. 4: Rückstandsuntersuchung.

EBRECHT, A.: Verbesserung des Hemmstofftestes durch Zusatz von Trimethoprim zum Nachweis von Sulfonamiden. - **Arch. Lebensmittelhyg.**, **33**; 109-115 (1982).

FORSCHNER, E., u. SEIDLER, M.: Alternativvorschläge zum Hemmstofftest. Rationalisierung und Absicherung. - **Fleischwirtsch.**, **56**; 1008-1013 (1976).

GUDDING, R.: An improved bacteriological method for the detection of sulfonamide residues in food. - **Acta Vet. Scand.**, **17**; 458-464 (1976).

LEVETZOW, R.: Untersuchungen auf Hemmstoffe im Rahmen der Bakteriologischen Fleischuntersuchung (BU). - Bundesgesundheitsblatt, 14; 15/16, 211-213 (1971).

PICHNARCIK, J., WENZEL, S., u. GISSKE, W.: Beitrag zur Methodik des Hemmstoffnachweises in Organen und Muskulatur von Schlachttieren. - **Arch. Lebensmittelhyg.**, **20**; 272-279 (1969).

Ordering Information

Product	Merck Cat. No.	Pack size
Test Agar pH 6.0 for the Inhibitor Test	1.10663.0500	500 g
Bacillus subtilis (BGA)-spore suspension	1.10649.0001	15 x 2 ml
Penicillin G potassium salt	CN Biosciences	
Streptomycin sulfate	CN Biosciences	

Quality control

Test strains	Inhibition zones in mm Ø			
	Gentamicin		Penicillin	Streptomycin
	10 µg	30 µg	10 IU	10 µ
Bacillus subtilis strain BGA (DSMZ 618)	20-28	22-30	36-48	19-27



Gentamicin 30µg



Penicillin 10IE