

## Product Information


# Peptidyl Arginine Deiminase type-2 (PAD2), human

Recombinant, expressed in *Escherichia coli***SAE0061**

## Product Description

Synonyms: Protein arginine iminohydrolase, PAD2, PADI2.

Peptidyl Arginine Deiminases (PADs) perform post translational deiminations of proteins. PADs are calcium-dependent enzymes that catalyze the conversion of L-arginine residues to L-citrulline. The reaction that is catalyzed is as follows:



This deimination provides another level of regulating protein function.

There are five mammalian PADs sub-types, which differ in substrate specificity and tissue distribution. PAD enzymes are highly homologous, with 50-60% sequence similarity.<sup>1</sup> PAD enzymes play important roles in gene regulation by citrullination of arginine residues on histones H3, H2A, and H4. Overexpression of these enzymes has been found in several diseases such as rheumatoid arthritis, Alzheimer's disease, multiple sclerosis, lupus, Parkinson's disease, and cancer.<sup>2,3</sup> PAD2 is widely expressed in the brain, secretory glands, and skeletal muscles. The enzyme was found to be responsible for hyper-citrullination of myelin basic protein, which is suspected to lead to multiple sclerosis.<sup>2</sup>

SAE0061 has been cited in the research literature.<sup>4</sup>

## Precautions and Disclaimer

This product is for R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

## Product

The product is supplied in 20 mM Tris (pH 7.5), 200 mM NaCl, 10% (w/v) glycerol, 10 mM 2-mercaptoethanol, 1 mM EDTA, 1 mM DTT, 10 mM CaCl<sub>2</sub>, 10 mM arginine, 30 mM maltose, and 0.6 mg/mL dodecyl-β-D maltoside.

Enzymatic activity: Activity is measured by a colorimetric method with the synthetic substrate benzoyl arginine ethyl ester (BAEE).<sup>5</sup>

Unit definition: One unit will produce 1 μmole of *N*-α-benzoylcitrulline ethyl ester from BAEE per hour at 55 °C at pH 7.2.

## Storage/Stability

Store the product at -20 °C. The product is stable for at least 2 years as supplied.

## Preparation Instructions

After first use, it is recommended to divide the enzyme into aliquots and immediately freeze at -20 °C. Avoid extended storage above freezing, as this will initially lead to an increase in enzymatic activity but may cause decreased stability. Avoid repeated freeze-thaw cycles. Do not store in a frost-free freezer.

## References

1. Mohanan, S., *et al.*, *Biochem. Res. Int.*, 895343 (doi: 10.1155/2012/895343) (2012).
2. Moscarello, M. A., *et al.*, *Neurochem. Res.*, 32(2), 251-256 (2007).
3. Bicker, K. L., and Thompson, P. R., *Biopolymers*, 99(2), 155-163 (2013).
4. Mahneva, O. *et al.*, *PLoS ONE*, 15(1), e0227822 (2020).
5. Takahara, H., *et al.*, *J. Biochem.*, 94(6), 1945-1953 (1983).

## Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

### Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://SigmaAldrich.com/techservice).

### Terms and Conditions of Use

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://SigmaAldrich.com/terms).

### Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://SigmaAldrich.com/offices).

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2017-2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.  
SAE0061pis Rev 12/23 AB, MP

The Merck logo, consisting of the word "MERCK" in a bold, red, sans-serif font.