


Product Information

Pyoverdines from *Pseudomonas fluorescens*

Catalog Number **P8124**

Storage Temperature –20 °C



CAS RN 8062-00-8

Synonym: Pseudobactins

Product Description

The product is a mixture containing primarily the following forms of pyoverdines:¹ succinic acid (MW = 1161) and succinamide (MW = 1160). The 2-hydroxyglutaramide (MW = 1190) form is a minor component.

Purity: >90% (HPLC)

Pyoverdines, also called pseudobactins, are fluorescent siderophores that have high-affinity for iron(III) ions (10^{32} M^{-1}) and are synthesized by fluorescent pseudomonads under iron-deficient growth conditions.¹ They are composed of three different structural parts: a dihydroxyquinoline chromophore, which confers color and fluorescence; a side chain, generally a dicarboxylic acid or a dicarboxylic amide; and a variable peptidic chain comprising 6 to 12 amino acids, some of them being unusual.^{1,3} The peptide chain contains two of the iron(III) binding sites, and the catecholate of the chromophore contains the third one. The peptide moiety is also involved in receptor recognition and binding.^{1,3} The size and amino acid composition of pyoverdines are unique to each species. Despite the strain specificity of pyoverdine recognition and uptake, many fluorescent pseudomonads are able to utilize pyoverdines produced by other strains.¹

Pyoverdines are not only iron chelators. The ferri-pyoverdine complex acts as a signaling molecule inducing the production of secreted virulence factors in *Pseudomonas aeruginosa*.⁴ Pyoverdines are able to prevent iron toxicity produced by iron overload in hepatocyte cultures⁵ and effectively scavenge hydroxyl and peroxyl radicals.⁶ Pyoverdines are also effective at acquiring iron from transferrin and lactoferrin.³ This siderophore is also involved in the suppression of pythium induced damping-off of tomato⁷ and promotion of growth in some higher-plants.⁸ As opposed to ferri-pyoverdines, which promote the growth of both *P. fluorescens* and *P. aeruginosa* in minimal medium, gallium-pyoverdines act as growth inhibitors of both strains,⁹ possibly due to gallium interference with bacterial iron metabolism.¹⁰

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

Soluble in water at 10 mg/ml.

Storage/Stability

Store at –20 °C. Under these conditions the product is stable for 2 years. A solution in water is stable for at least 4 months at 2–8 °C.

Related Products

Pyoverdines - Fe complex (Catalog Number P8374)
Pyoverdines - Ga(III) complex (Catalog Number P8249)

References

1. Linget, C., et al., Bacterial siderophores: the structures of the pyoverdins of *Pseudomonas fluorescens* ATCC 13525. *Tetrahedron lett.*, **33**, 1737-1740 (1992).
2. Fuchs, R., et al., Siderotyping - a powerful tool for the characterization of pyoverdines. *Curr. Top. Med. Chem.*, **1**, 31-57 (2001).
3. Poole, K., and McKay, G.A., Iron acquisition and its control in *Pseudomonas aeruginosa*: many roads lead to Rome. *Front. Biosci.*, **8**, 661-686 (2003).
4. Beare, P.A., et al., Siderophore-mediated cell signaling in *Pseudomonas aeruginosa*: divergent pathways regulate virulence factor production and siderophores receptor synthesis. *Mol. Microbiol.*, **47**, 195-207 (2003).
5. Jego, P., et al., Inhibition of iron overload toxicity in rat hepatocyte cultures by pyoverdin Pf, the siderophores of *Pseudomonas fluorescens*. *Biochem. Pharmacol.*, **43**, 1275-1280 (1992).
6. Morel, I., et al., Antioxidant and free radical scavenging activities of the iron chelators pyoverdin and hydroxypyrid-4-ones in iron-loaded hepatocyte cultures: comparison of their mechanism of protection with that of desferrioxamine. *Free Rad. Biol. Med.*, **13**, 499-508 (1992).
7. Buysens, S., et al., Involvement of pyochelin and pyoverdin in suppression of pythium-induced damping-off of tomato by *Pseudomonas aeruginosa* 7NSK2. *Appl. Environ. Microbiol.*, **62**, 865-871 (1996).
8. Boruah, H.P., and Kumar, B.S., Biological activity of secondary metabolites produced by a strain of *Pseudomonas fluorescens*. *Folia Microbiol.*, **4**, 359-363 (2002).
9. Schlosser-Silverman, E., et al., Biological applications of pyoverdines and their complexes with iron and gallium. Society for Industrial Microbiology Annual meeting, Baltimore, USA (2006).
10. Olakanmi, O., et al., Gallium disrupts iron metabolism of mycobacteria residing within human macrophages. *Infect. Immun.*, **10**, 5619-5627 (2000).

KAA,ES,MAM 10/06-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.