

3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

# **ProductInformation**

## Copper(II) sulfate pentahydrate

Product Number **C 8027** Store at Room Temperature

## **Product Description**

Molecular Formula: CuSO<sub>4</sub> • 5H<sub>2</sub>O

Molecular Weight: 249.7 CAS Number: 7758-99-8

Synonym: cupric sulfate pentahydrate

This product is cell culture tested (0.02  $\mu$ g/ml) and is appropriate for use in cell culture applications.

Copper sulfate is a reagent that is used in many largescale applications. Applications include textile dyeing, the preparation of azo dyes, wood preservation, and the tanning of leather. Copper sulfate pentahydrate occurs in nature as the mineral chalcanthite.<sup>1</sup>

Copper sulfate is frequently utilized to oxidize lipoproteins in the context of biological oxidative stress. <sup>2,3</sup> It is also used in studies of plants and plant pathogens. <sup>4,5</sup>

A protocol for the isolation of recombinant yeast 6-phosphofructo-2-kinase that uses copper sulfate has been published.<sup>6</sup> A procedure has been reported for the purification of dye- and metal ion-binding proteins in a polyvinylpyrrolidone-based aqueous two-phase system.<sup>7</sup>

### **Precautions and Disclaimer**

For Laboratory Use Only. Not for drug, household or other uses.

## **Preparation Instructions**

This product is soluble in water (100 mg/ml), yielding a clear blue solution. It is also soluble in methanol and glycerol, and slightly soluble in ethanol.<sup>1</sup>

#### References

- 1. The Merck Index, 12th ed., Entry# 2722.
- Chang, M. Y., et al., Oxidized LDL bind to nonproteoglycan components of smooth muscle extracellular matrices. J. Lipid Res., 42(5), 824-833 (2001).
- Nguyen-Khoa, T., et al., Oxidized low-density lipoprotein induces macrophage respiratory burst via its protein moiety: A novel pathway in atherogenesis? Biochem. Biophys. Res. Commun., 263(3), 804-809 (1999).
- Grey, B. E., and Steck, T. R., The viable but nonculturable state of *Ralstonia solanacearum* may be involved in long-term survival and plant infection. Appl. Environ. Microbiol., 67(9), 3866-3872 (2001).
- Muller, S. L., et al., Effects of copper sulfate on Typha latifolia seed germination and early seedling growth in aqueous and sediment exposures. Arch. Environ. Contam. Toxicol., 40(2), 192-197 (2001).
- Dihazi, H., et al., One-step purification of recombinant yeast 6-phosphofructo-2-kinase after the identification of contaminants by MALDI-TOF MS. Protein Expr. Purif., 21(1), 201-209 (2001).
- 7. Fernandes, S., et al., Affinity extraction of dye- and metal ion-binding proteins in polyvinylpyrrolidone-based aqueous two-phase system. Protein Expr. Purif., **24(3)**, 460-469 (2002).

GCY/RXR 4/03