



3050 Spruce Street  
Saint Louis, Missouri 63103 USA  
Telephone 800-325-5832 • (314) 771-5765  
Fax (314) 286-7828  
email: techserv@sial.com  
sigma-aldrich.com

## Product Information

### Anti-CENP-A

produced in rabbit, affinity isolated antibody

Catalog Number **C6867**

#### Product Description

Anti-CENP-A is produced in rabbit using a synthetic peptide corresponding to amino acids 32-49 located near the N-terminus of human CENP-A, conjugated to KLH, as immunogen. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-CENP-A recognizes human CENP-A. 17 kDa, by immunoblotting. Staining of the CENP-A band in immunoblotting is specifically inhibited with the immunizing peptide.

The centromere is an essential chromosomal component that is required for proper segregation of chromosomes during mitosis and meiosis. The kinetochore is a large DNA-multiprotein complex that mediates attachment of microtubules to centromere during mitosis. It comprises both constitutive proteins that are present at the centromere throughout the cell cycle and transient proteins that are present at various stages. The centromere proteins (CENP) CENP-A, CENP-B and CENP-C are constitutive proteins of the kinetochore. CENP-A (also known as CenH3) is a 17-kDa histone H3-like protein involved in centromeric nucleosome formation.<sup>1,2</sup> CENP-A is specifically incorporated in direct contact with DNA in the region of the inner kinetochore plate of active centromeres.<sup>3-5</sup> CENP-A has a fold domain similar to histone H3. The C-terminal of CENP-A is 62% identical to histone H3 and is required for targeting to the centromere. In contrast, there is no similarity between the N-terminal (1-47) regions of CENP-A and histone H3. CENP-A has been shown to replace histone H3 in nucleosome reconstitution *in vitro*.<sup>6</sup> It has been shown that CENP-A is phosphorylated at Ser<sup>7</sup> by both Aurora-A and Aurora-B protein kinases in early mitosis.<sup>7-9</sup> CENP-A phosphorylation is required for proper kinetochore attachment to microtubules and chromosome alignment during mitosis. CENP-A is phosphorylated in early prophase after completion of histone H3 phosphorylation and is dephosphorylated in early anaphase, before completion of histone H3 dephosphorylation. The temporal differences at G<sub>2</sub>M between CENP-A and histone H3 phosphorylation suggest that although they

share a similar phosphorylation site, they serve different functions. Knock-out of the CENP-A gene in mouse is lethal.<sup>10</sup> Affected embryos showed severe mitotic problems, including micronuclei and macronuclei formation, nuclear bridging and blebbing, chromatin fragmentation and hypercondensation. CENP-A depletion results in diffuse CENP-B foci, absence of discrete CENP-C signal on centromeres, and dispersion of CENP-B and CENP-C throughout the nucleus. It has been suggested that CENP-A is essential for kinetochore targeting of CENP-C and plays an early role in organizing centromeric chromatin at interphase.

#### Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Antibody concentration: ~2 mg/mL

#### Precautions and Disclaimer

Due to the sodium azide content, a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazards and safe handling practices.

#### Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

#### Product Profile

Immunoblotting: a working concentration of 1-2 µg/mL is recommended using HEK-293 cells expressing human CENP-A.

**Note:** In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

## References

1. Palmer, D.K., et al., *Proc. Natl. Acad. Sci. USA*, **88**, 3734-3738 (1991).
2. Sullivan, K.F., et al., *J. Cell. Biol.*, **127**, 581-592 (1994).
3. Warburton, P.E., et al., *Curr. Biol.*, **7**, 901-904 (1997).
4. Shelby, R.D., et al., *J. Cell. Biol.*, **136**, 501-513 (1997).
5. Van Hooser, A.A., et al., *J. Cell. Sci.*, **114**, 3529-3542 (2001).
6. Yoda, K., et al., *Proc. Natl. Acad. Sci. USA*, **97**, 7266-7271 (2000).
7. Zeitlin, S.G., et al., *J. Cell. Sci.*, **114**, 653-661 (2000).
8. Zeitlin, S.G., et al., *J. Cell. Biol.*, **155**, 1147-1157 (2001).
9. Kunitoku, N., et al., *Dev. Cell*, **5**, 853-864 (2003).
10. Howman, E.V., et al., *Proc. Natl. Acad. Sci. USA*, **97**, 1148-1153 (2000).

ER,KAA,PHC 08/06-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.