

## Product Information

# Hirudin

From leeches

**H7016, H7380**

Storage Temperature: 2–8 °C

CAS RN 8001-27-2

## Product Description

The anticoagulant, hirudin, is the most potent natural inhibitor of both soluble and clot-bound thrombin. It binds thrombin with high affinity, the hirudin- $\alpha$ -thrombin complex has a dissociation constant of  $\sim 1 \times 10^{-11}$  M.<sup>1,2</sup> Hirudin covers more than 20% of the surface area of thrombin, occluding both the active site and exosite I (fibrinogen and PAR recognition site). This coverage blocks thrombus growth and platelet activation. Hirudin is not metabolized in the bloodstream of humans and is eliminated unchanged via kidney filtration.

Hirudin is a ~7 kDa acidic protein containing 65 amino acid residues. It contains a sulfated tyrosyl residue (Tyr<sup>63</sup>), three disulfide bridges, and a high proportion of dicarboxylic acids.<sup>2,3</sup> Hirudin is not glycosylated and lacks tryptophan, arginine, and methionine residues. At least 20 isoforms have been identified and sequenced.<sup>3</sup>

Isoelectric point (pI): 3.5–4.0

Hirudin is produced in the salivary glands of the leech *Hirudo verbana*.<sup>4</sup> Hungry leeches provide a richer source of hirudin.<sup>4</sup> The method of preparation is a modification of published procedures.<sup>1,4</sup> Both products are supplied as lyophilized powders.

Specific Activity:

- H7016,  $\geq 1,500$  antithrombin units/mg protein
- H7389, 300–1,500 antithrombin units/mg protein

One antithrombin unit (ATU) will neutralize one NIH unit of thrombin at 37 °C, based on direct comparison to an NIH thrombin reference standard.

## Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

## Preparation Instructions

Hirudin is soluble in water. The literature cites the use of "Dilution Fluid II" (35.7 mM acetic acid, 35.7 mM sodium diethyl barbiturate, 0.85% NaCl, 1% bovine serum albumin, and 0.5% PEG, pH 7.4) to dissolve hirudin (500 ATU/mL) and thrombin.<sup>5</sup>

Hirudin is soluble in pyridine, but practically insoluble in alcohol, ether, acetone, or benzene.

## Storage/Stability

Store the products at 2–8 °C.

## References

1. Markwardt, F., Methods in Enzymology, 19, 924–932 (1970).
2. Dodt, J., et al., FEBS Letters, 165, 180–184 (1984), citing Markwardt, F., and Walsmann, P., Hoppe-Seyler's Z. Physiol. Chem., 312, 85–98 (1958).
3. Stone, S.R., and Maraganore, J.M., Methods in Enzymology, 223, 312 (1993).
4. Bagdy, D., et al., Methods in Enzymology, 45, 669–678 (1976).
5. Loison, G., et al., Bio/Technology, 6, 72 (1988).

---

## Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

### Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://SigmaAldrich.com/techservice).

### Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://SigmaAldrich.com/terms).

### Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://SigmaAldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2008-2024 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.  
H7380pis Rev 05/24

**Millipore  
SIGMA**