

Product Information

Nuclease Micrococcal

From *Staphylococcus aureus*

N3755

Storage Temperature –20 °C

Product Description

Enzyme Commission (EC) Number: 3.1.31.1

CAS Number: 9013-53-0

Molecular Weight: 16,807 (based on sequence)¹

Extension coefficient: 0.935

This enzyme has an absolute need for Ca^{2+} for activity. The pH optimum varies according to Ca^{2+} concentration.² The optimal pH for RNase and DNase activities is between 9 and 10, depending on the Ca^{2+} concentration. At higher pH values, less Ca^{2+} is required. The inhibitory effect of high Ca^{2+} concentrations is more pronounced at higher pH values.³ Mg^{2+} cannot replace Ca^{2+} in activating the enzyme.⁴

This enzyme will cleave DNA and RNA to leave 3'-nucleotides.

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

Preparation Instructions

Reconstitute to a concentration of 1 unit per 5 µL with water.

Storage/Stability

After reconstitution with water, aliquots may be frozen by immersion in a dry ice/alcohol bath and then stored at –20 °C. It is suggested that stability is improved if the product is dissolved in 0.1% BSA to minimize adsorption to container walls. Aliquots may also be re-lyophilized.

Protocol

For Nuclease Treatment

1. Transfer 500 µL of reconstituted lysate to a separate vial.
2. Add 5 µL of micrococcal nuclease solution (1 unit per 5 µL).
3. Start the reaction by adding 5 µL of 0.1 M CaCl_2 solution.
4. Mix gently and constantly for 2 minutes in a 28 °C water bath.
5. At the end of 2 minutes, add 10 µL of 0.1 M EGTA to stop the reaction and place the vial in an ice bath. Wait two minutes. Sample is ready to use.

References

1. Taniuchi, H., Anfinsen, C. B., & Sodja, A. (1967). The Amino Acid Sequence of an Extracellular Nuclease of *Staphylococcus aureus*: III. COMPLETE AMINO ACID SEQUENCE. *Journal of Biological Chemistry*, 242(20), 4752-4758.
2. Heins, J. N., Suriano, J. R., Taniuchi, H., & Anfinsen, C. B. (1967). Characterization of a nuclease produced by *Staphylococcus aureus*. *Journal of Biological Chemistry*, 242(5), 1016-1020.
3. Cuatrecasas, P., Fuchs, S., & Anfinsen, C. B. (1967). Catalytic properties and specificity of the extracellular nuclease of *Staphylococcus aureus*. *Journal of Biological Chemistry*, 242(7), 1541-1547.
4. Reddi, K. K. (1967). [33] Micrococcal nuclease. In *Methods in enzymology* (Vol. 12, pp. 257-262). Academic Press.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck operates
as MilliporeSigma in the U.S. and Canada.

Merck, and Sigma-Aldrich are trademarks of Merck or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2011-2025 Merck and/or its affiliates. All Rights Reserved.
23276664 Rev 11/25

