

SIGMA QUALITY CONTROL TEST PROCEDURE**Enzymatic Assay of GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE¹
(EC 1.2.1.12)****PRINCIPLE:**

Abbreviations used:

3-PGA = 3-Phosphoglyceric Acid

ATP = Adenosine 5'-Triphosphate

3-PGK = 3-Phosphoglyceric Phosphokinase

ADP = Adenosine 5'-Diphosphate

NADH = Nicotinamide Adenine Dinucleotide, Reduced Form

GAPDH = Glyceraldehyde-3-Phosphate Dehydrogenase

G-3-P = Glyceraldehyde 3-Phosphate

NAD = Nicotinamide Adenine Dinucleotide, Oxidized Form

P_i = Inorganic Phosphate**CONDITIONS:** T = 25°C, pH = 7.6, A_{340nm}, Light path = 1 cm**METHOD:** Continuous Spectrophotometric Rate Determination**REAGENTS:**

- A. 100 mM Triethanolamine Buffer, pH 7.6 at 25°C
(Prepare 100 ml in deionized water using Triethanolamine Hydrochloride, Sigma Prod. No. T-1502. Adjust to pH 7.6 at 25°C with 1 M NaOH. **Prepare fresh.**)
- B. 100 mM 3-Phosphoglyceric Acid Solution (3-PGA)
(Prepare 2 ml in deionized water using D(-)-3-Phosphoglyceric Acid, tri(cyclohexylammonium) Salt, Sigma Prod. No. P-8752.)
- C. 200 mM L-Cysteine HCl Solution (Cys)
(Prepare 1 ml in deionized water using L-Cysteine Hydrochloride, Monohydrate, Sigma Prod. No. C-7880. Neutralize the solution by adding solid Sodium Bicarbonate, Sigma Prod. No. S-8875. **PREPARE FRESH.**)

Enzymatic Assay of GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE¹
(EC 1.2.1.12)

REAGENTS: (continued)

- D. 100 mM Magnesium Sulfate Solution (MgSO₄)
(Prepare 10 ml in deionized water using Magnesium Sulfate, Heptahydrate, Sigma Prod. No. M-1880.)
- E. 7.0 mM β -Nicotinamide Adenine Dinucleotide, Reduced Form Solution (β -NADH)
(Prepare 1 ml in deionized water using β -Nicotinamide Adenine Dinucleotide, Reduced Form, Disodium Salt, Sigma Prod. No. N-8129 or dissolve the contents of one 5 mg vial of β -Nicotinamide Adenine Dinucleotide, Reduced Form, Disodium Salt, Sigma Stock No. 340-105, in the appropriate volume of deionized water. **PREPARE FRESH.**)
- F. 34 mM Adenosine 5'-Triphosphate Solution (ATP)
(Prepare 1 ml in deionized water using Adenosine 5'-Triphosphate, Disodium Salt, Sigma Prod. No. A-5394. **PREPARE FRESH.**)
- G. 3-Phosphoglyceric Phosphokinase Enzyme Solution (3-PGK)
(Immediately before use, prepare a solution containing 200 units/ml in cold deionized water using 3-Phosphoglyceric Phosphokinase, Sigma Prod. No. P-7634.)
- H. Glyceraldehyde-3-Phosphate Dehydrogenase Enzyme Solution (GAPDH)
(Immediately before use, prepare a solution containing 0.3 - 0.6 unit/ml of Glyceraldehyde-3-Phosphate Dehydrogenase in cold Reagent A.)

PROCEDURE:

Pipette (in milliliters) the following reagents into suitable cuvettes:

	<u>Test</u>	<u>Blank</u>
Reagent A (Buffer)	2.40	2.50
Reagent B (3-PGA)	0.20	0.20
Reagent C (Cys)	0.05	0.05
Reagent D (MgSO ₄)	0.05	0.05
Reagent E (β -NADH)	0.05	0.05
Reagent F (ATP)	0.10	0.10
Reagent G (3-PGK)	0.05	0.05

Enzymatic Assay of Glyceraldehyde-3-PHOSPHATE DEHYDROGENASE¹ (EC 1.2.1.12)

PROCEDURE: (continued)

Mix by inversion and equilibrate to 25°C. Monitor the $A_{340\text{nm}}$ until constant, using a suitably thermostatted spectrophotometer. Then add:

	<u>Test</u>	<u>Blank</u>
Reaction H (Enzyme Solution)	0.10	-----

Immediately mix by inversion and record the decrease in $A_{340\text{nm}}$ for approximately 5 minutes. Obtain the $\Delta A_{340\text{nm}}/\text{minute}$ using the maximum linear rate for both the Test and Blank.

CALCULATIONS:

$$\text{Units/ml enzyme} = \frac{(\Delta A_{340\text{nm}}/\text{min Test} - \Delta A_{340\text{nm}}/\text{min Blank})(3)(\text{df})}{(0.1)(6.22)}$$

3 = Volume (in milliliters) of assay

df = Dilution factor

0.1 = Volume (in milliliter) of enzyme used

6.22 = Millimolar extinction coefficient of α -NADH at 340 nm

$$\text{Units/mg solid} = \frac{\text{units/ml enzyme}}{\text{mg solid/ml enzyme}}$$

$$\text{Units/mg protein} = \frac{\text{units/ml enzyme}}{\text{mg protein/ml enzyme}}$$

UNIT DEFINITION:

One unit will reduce 1.0 μ mole of 3-phosphoglycerate to D-glyceraldehyde-3-phosphate per minute in a coupled system with 3-phosphoglyceric phosphokinase at pH 7.6 at 25°C.

FINAL ASSAY CONCENTRATION:

In a 3.00 ml reaction mix, the final concentrations are 83 mM triethanolamine, 6.7 mM 3-phosphoglyceric acid, 3 mM L-cysteine, 2 mM magnesium sulfate, 0.1 mM α -NADH, 1.1 mM ATP, 10 units 3-phosphoglyceric phosphokinase and 0.03 - 0.06 unit glyceraldehyde-3-phosphate dehydrogenase.

Enzymatic Assay of GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE¹
(EC 1.2.1.12)

NOTES:

1. Not to be used to assay activity of Glyceraldehyde-3-Phosphate Dehydrogenase, from *Bacillus Stearothermophilus*, Sigma Prod. No. G-5892.
2. 3-Phosphoglyceric Phosphokinase unit definition: One unit will convert 1.0 μ mole of 1,3-diphosphoglycerate to 3-phosphoglycerate per minute at pH 6.9 at 25°C.
3. All product and stock numbers, unless otherwise indicated, are Sigma product and stock numbers.

Sigma warrants that the above procedure information is currently utilized at Sigma and that all Sigma-Aldrich, Inc. products conform to the information in this and other Sigma-Aldrich, Inc. publications. Purchaser must determine the suitability of the information and product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.