

Product Information

Mpro, 3CL Protease from coronavirus SARS-COV2

Recombinant protein, lyophilized powder

SAE0172

Product Description

Maturation of the SARS-CoV2 virus depends on cleavage of the overlapping large polyproteins 1a and 1ab by two viral proteases:

- Mpro (main protease)
- PLpro (Papain-like protease)

Proteolytic cleavage by Mpro of the 1ab polyprotein occurs at 11 sites:

- 7 sites within the 1a polyprotein
- 4 sites within the 1ab polyprotein

This results in maturation of a total of 16 viral non-structural proteins (NSP).^{1,2}

Mpro protease forms a functional homodimer. Both the *N*-terminus and the *C*-terminus of Mpro have been shown to be critical for dimer formation and for enzyme function.²

The Mpro protease is an ideal target for antiviral drug design due to its high conservation between different coronavirus strains and absence of functional analogs in the human proteome.^{2,3} It is noteworthy that Mpro protease from SARS-CoV1 and SARS-CoV2 are functionally identical.^{4,5}

This fully active Mpro protease product contains the complete sequence of Mpro protease (Accession: YP_009725301.1) without any additional tags. It is provided as a lyophilized powder for increased stability.

This product has been used in a Surface Plasmon Resonance (SPR) binding assay.⁶

Reagent

The product is supplied lyophilized from 20 mM HEPES (pH 7.3), 2.5% Trehalose, and 0.05% TWEEN® 20.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

The product retains activity for at least 2 years when stored lyophilized at -20 °C.

Preparation Instructions

It is suggested to reconstitute the product in water or in 10% aqueous glycerol solution at a protein concentration of 1-2 mg/mL. Aliquot the protein solution and store the solution aliquots at -20 °C. Avoid freeze-thaw cycles.

Procedure

Activity measurement using fluorogenic substrate peptide (Cat. No. SAE0180)

- Prepare the stock solution of the fluorogenic peptide in DMSO at a concentration of 20 mg/mL.
- Depending on the hardware set-up, the fluorogenic peptide can be used at working concentrations of 10-100 µg/mL.

Sample Protocol

1. Prepare the substrate solution at a final concentration of 10 µg/mL in 25 mM HEPES, pH 7.0.
2. Set up the fluorimeter (fluorescent plate reader) for reading at an excitation (λ_{ex}) of 400 nm, and emission (λ_{em}) at 505 nm.
3. Use 200 µL of substrate solution per well for the 96-well plate format.
4. Add 2-20 µg/mL Mpro.
5. Read kinetics for 5-60 minutes.




Figure 1. Mpro enzyme kinetics performed with the fluorogenic substrate SAE0180.

- Substrate concentration: 10 µg/mL
- Mpro enzyme concentrations: 0, 2.5 and 5 µg/mL

Experiments with Mpro at 2.5 and 5 µg/mL are performed in triplicate.

References

1. Fan, K. *et al.*, *J. Biol. Chem.*, **279(3)**, 1637-1642 (2004).
2. Xia, B., and Kang, X., *Protein Cell.*, **2(4)**, 282-290 (2011).
3. Yang, H. *et al.*, *Curr. Pharm. Des.*, **12(35)**, 4573-4590 (2006).
4. Goyal, B., and Goyal, D., *ACS Comb Sci.*, **22(6)**, 297-305 (2020).
5. Zhang, L. *et al.*, *Science*, **368(6489)**, 409-412 (2020).
6. Li, H. *et al.*, *J. Agric. Food Chem.*, **69(41)**, 12197-12208 (2021).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

SAE0172pis Rev 02/22 DK,DT,GCY

MERCK