

## Product Information

## Trypsin-Agarose

Buffered aqueous suspension, enzyme from bovine pancreas

**T1763**

### Product Description

Storage Temperature: 2-8 °C

Trypsin is a member of the serine protease family. The active site amino acid residues of trypsin include His<sup>46</sup> and Ser<sup>183,2-4</sup>. Trypsin is a single chain polypeptide of 223 amino acid residues. Trypsin is produced by the removal of the N-terminal hexapeptide from trypsinogen which is cleaved at the Lys<sup>6</sup>-Ile<sup>7</sup> peptide bond. Trypsin contains 6 disulfide bridges as cross-linkages. This native form of trypsin is referred to as β-trypsin. Autolysis of β-trypsin (which is cleaved at Lys<sup>131</sup>-Ser<sup>132</sup>) results in α-trypsin, which is held together by disulfide bridges.

Trypsin cleaves peptides on the C-terminal side of lysine and arginine amino acid residues. The rate of hydrolysis is slower if an acidic residue is on either side of the cleavage site. No cleavage occurs if a proline residue is on the carboxyl side of the cleavage site. The pH optimum of trypsin is 7-9.<sup>6</sup>

This trypsin-agarose product is prepared by the immobilization of trypsin, originally isolated from bovine pancreas, to activated crosslinked beaded agarose. Several references<sup>7-16</sup> have cited use of this product in their research applications.

### Precautions and Disclaimer

This product is for R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

### Product

This trypsin-agarose product is sold as a suspension in ~10 mM acetic acid, pH 3.2.

### Preparation Instructions

General instructions for re-suspension of our enzyme-agarose conjugates include the following steps.

- Suspend the lyophilized enzyme-agarose to 5-10 mg solid/mL water.
- Allow brief hydration of the lyophilized powder.
- Filter and wash the rehydrated enzyme-agarose product several times with either water or your intended buffer.
- Re-suspend the enzyme-agarose in your intended buffer. The product is now ready for use.

### Storage/Stability

For re-use of our enzyme-agarose conjugates, the following steps may be used as a general guide:

- Wash the enzyme-agarose with water and/or buffer until it is free of substrates. This specific enzyme-agarose conjugate may then be resuspended in a fresh stock of the parent buffer system.
- For long-term storage, enzyme-agarose products may be re-converted to their dry form, if desired, as follows:
  - Wash the enzyme-agarose with the buffer of choice.
  - Drain excess buffer.
  - Dry the enzyme-agarose in a vacuum desiccator.
  - Store the freshly lyophilized enzyme-agarose at 2-8 °C.

## References

1. Cunningham, L. W., Jr., *J. Biol. Chem.*, **211(1)**, 13-19 (1954).
2. Walsh, K. A., *Meth. Enzymol.*, **19**, 41-63 (1970).
3. Keil, B., in *The Enzymes*, 3rd ed., Vol. III (Boyer, P. D., ed.). Academic Press (New York), pp. 250-275 (1971).
4. Shaw, E., et al., *Biochemistry*, **4(10)**, 2219-2224 (1965).
5. Buck, F. F. et al., *Arch. Biochem. Biophys.*, **97**, 417-424 (1962).
6. Sipos, T., and Merkel, J. R., *Biochemistry*, **9(14)**, 2766-2775 (1970).
7. Bolte, G. et al., *Clin. Chem. Acta*, **247(1-2)**, 59-70 (1996).
8. Fraser, J. S. et al., *Gut*, **52(12)**, 1698-1702 (2003).
9. Hobson, J. P. et al., *J. Biol. Chem.*, **279(45)**, 46981-46984 (2004).
10. Bracken, S. C. et al., *Aliment. Pharmacol. Ther.*, **23(9)**, 1307-1314 (2006).
11. Bøgh, K. L. et al., *Clin. Exp. Allergy*, **39(10)**, 1611-1621 (2009).
12. Zevallos, V. F. et al., *Am. J. Clin. Nutr.*, **96(2)**, 337-344 (2012).
13. Wahab, W. A. et al., *Int. J. Exp. Pathol.*, **97(4)**, 303-309 (2016).
14. Han, B. et al., *ACS Nano*, **13(10)**, 11320-11333 (2019).
15. Jiang, J. et al., *Nat. Chem.*, **15(4)**, 578-586 (2023).
16. Olías, R. et al., *Food Res. Int.*, **169**, 112825 (2023).

## Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

### Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

### Terms and Conditions of Use

Warranty, use restrictions, and other conditions of sale may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

### Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, Trizma and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 1999-2023 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

T1763pis Rev 10/23 GCY,MAM

**MILLIPORE  
SIGMA**