

3050 Spruce Street, St. Louis, MO 63103 USA
Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757
email: techservice@sial.com sigma-aldrich.com

Product Information

Casein solution, 5% from bovine milk

Catalog Number **C4765** Storage Temperature 2–8 °C

CAS RN (for casein) 9000-71-9

Product Description

This hydrolyzed and dephosphorylated casein product is a sterile filtered solution in purified water. It is suitable for use as a substrate for 3′,5′-cyclic-AMP dependent protein kinase.

Casein is a phosphoprotein found in milk. This protein has numerous experimental applications, including use as a blocking agent in immunochemistry, recovery of enzyme activity from SDS extracted samples, and as a substrate for protease and kinase assays. The major casein subunits may be distinguished by electrophoresis, and are designated as α -, β -, γ -, and κ -caseins in order of decreasing mobility at pH 7.0. The approximate casein composition of milk is (g/L): α -s1 (12–15); α -s2 (3–4); β (9–11); and κ (2–4).

The casein subunits vary primarily in molecular mass, isoelectric point, and level of phosphorylation. The following table lists these differences.^{2,3}

Subunit	MW (kDa)	pl	Phosphates/ mole	E ^{1%} (280 nm)
α-s1	22-23.7	4.2-4.7	8–10	10.0–10.1
α-s2	25	-	10–13	_
β	24	4.6–5.1	4–5	4.5-4.7
κ	19	4.1–5.8	1	10.5

The nomenclature for proteins in bovine milk has been published.²

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

References

- 1. The Merck Index 11th ed., Entry# 1892.
- Eigel, W.N. et al., J. Dairy Sci., 67(8), 1599-1631 (1984).
- 3. Modler, H.W., J. Dairy Sci., **68(9)**, 2195-2205 (1985).

SG,GY,VR,MWM,NSB,RBG,MAM 04/15-1