

Product Information

Activin A Human

recombinant, expressed in HEK 293 cells, HumanKine®, suitable for cell culture

H4666

Storage temperature -20 °C

Description

HumanKine® Activin A is expressed as a glycosylated disulfide linked 25 kDa homodimer in HEK 293 cells.¹ Glycosylation contributes to stability in cell growth media and other applications. Production in human 293 cells offers authentic glycosylation. Glycosylation contributes to stability in cell growth media and other applications.

Activin A (β A- β A) is secreted by Sertoli cells in the testis² and granulosa cells in the ovary. In early studies, this peptide was thought to be an inhibin and not recognized as a unique compound.^{3,4} Activins and inhibins are members of the TGF- β superfamily due to amino acid homology with respect to the conservation of 7 of the 9 cysteine residues common to all TGF- β forms.⁴ Activins are homodimers or heterodimers of the various β subunit isoforms, while inhibins are heterodimers of a unique α subunit and one of the various β subunits.⁵ Five β subunits have been cloned (mammalian β A, β B, β C, β E, and Xenopus β D). The activin/inhibin nomenclature reflects the subunit composition of the proteins: activin A (β A- β A), activin B (β B- β B), activin AB (β B- β A), inhibin A (α - β A), and inhibin B (α - β B). The mature human β A subunit is 100% identical to mouse β A. Cells known to express the β A chain include fibroblasts, endothelial cells, vascular smooth muscle cells, hepatocytes, macrophages, keratinocytes, osteoclasts, prostatic epithelium, neurons, chondrocytes, osteoblasts, Leydig cells, bone marrow monocytes, Sertoli cells, and ovarian granulosa cells.

Activins have a wide range of biological activities including mesoderm induction,^{6,7} bone remodeling, neural cell differentiation, hematopoiesis, and reproductive physiology. Activin A is involved in growth and differentiation of several tissues from different species.^{2-3,7-8} This protein also plays a key role in the production and regulation of hormones such as FSH, LH, GnRH, and ACTH. Activin influences erythropoiesis and the potentiation of erythroid colony

formation, oxytocin secretion, paracrine, and autocrine regulation.³

Similar to other TGF- β family members, activins exert their biological activities through the effects of the heterodimeric complex composed of two membrane spanning serine-threonine kinases designated type I and type II receptors.⁹ Activin type I and type II receptors are distinguished by the level of sequence homology of their kinase domains, and other structural and functional features. To date, seven type I and five type II activin receptors have been cloned from mammals, including activin receptor IA, activin receptor IIA, activin receptor IB, and activin receptor IIB. In addition, two splice variants of activin receptor IIA and five splice variants of activin receptor IIB have been reported.¹⁰ Activin binds directly to activin receptor type II, this complex then associates with activin receptor type I and initiates signal transduction.¹¹

This product is lyophilized from a PBS solution.

EC₅₀: \leq 5.0 ng/mL

The specific activity is determined by the dose-dependent inhibition of proliferation of the MPC-11 cell line (mouse plasmocytoma cell line).

Purity: \geq 95% (SDS-PAGE)

Endotoxin level: \leq 1 EU/ μ g

Preparation Instructions

Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile PBS containing 0.1% endotoxin-free, recombinant human serum albumin

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Store the product at -20 °C. The lyophilized product remains active for one year at -20 °C. Upon reconstitution, the cytokine can be stored at 2-8 °C for short term only, or at -20 °C to -80 °C in aliquots for long term. Avoid repeated freeze-thaw cycles.

References

1. Mason, A.J., et al., Structure of two human ovarian inhibins. *Biochem. Biophys. Res. Commun.*, **135**, 957-964 (1986).
2. de Winter, J., et al., Activin is produced by rat Sertoli cells in vitro and can act as an autocrine regulator of Sertoli cell function. *Endocrinology*, **132**, 975-982 (1993).
3. Sporn, M.B., and Roberts, A.B., eds., *Peptide Growth Factors and Their Receptors*, Vol. II, Springer-Verlag (Heidelberg, Germany: 1991) pp 217-235.
4. De Jong, F., et al., Effects of factors from ovarian follicular fluid and Sertoli cell culture medium on *in vivo* and *in vitro* release of pituitary gonadotrophins in the rat: an evaluation of systems for the assay of inhibin. *J. Reprod. Fertil.*, **26**, 47-59 (1979).
5. Vale, W. et al., The Inhibin/Activin Family of Hormones and Growth Factors. in *Peptide Growth factors and their Receptors*. Sporn, M., and Roberts, A., eds., Springer-Verlag, (Berlin, Germany: 1990) pp. 211-248.
6. Smith, J., et al., Expression of a *Xenopus* homolog of Brachyury (T) is an immediate-early response to mesoderm induction. *Cell*, **67**, 79-87 (1991).
7. Ying, S.Y., et al., Activins and activin receptors in cell growth. *Proc. Soc. Exp. Biol. Med.*, **214**, 114-122 (1997).
8. Woodruff, T.K., Regulation of cellular and system function by activin. *Biochem. Pharmacol.*, **55**, 953-963 (1998).
9. ten Dijke, P., et al., Activin receptor-like kinases: a novel subclass of cell-surface receptors with predicted serine/threonine kinase activity. *Oncogene*, **8**, 2879-2887 (1993).
10. Shoji, H., et al., Identification and Characterization of a PDZ Protein That Interacts with Activin Type II Receptors. *J. Biol. Chem.*, **275**, 5485-5492 (2000).
11. Attisano, L., et al., Identification of human activin and TGF beta type I receptors that form heteromeric kinase complexes with type II receptors. *Cell*, **75**, 671-680 (1993).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

**MILLIPORE
SIGMA**