Sigma-Aldrich_®

Product Information Sheet

Acetyl Coenzyme A Trisodium Salt

A2056

Product Description

Molecular Formula: C₂₃H₃₅N₇O₁₇P₃SNa₃ (for free acid)

Molecular Weight: 875.51 (for free acid)

E^M: 16,400 (water, 260 nm)¹

E^M: 15,400 (0.1 M PO4 Buffer, pH 7, 259 nm)

EM: 8,700 (water, 232 nm)1

This product, A2056, is prepared enzymatically by reacting coenzyme A (CoA) with acetyl phosphate and phosphotransacetylase. It is purified by ion exchange chromatography. Several methods of preparation and methods for the determination of Acetyl CoA and other CoA derivatives have been published.²⁻⁴

Coenzymes comprise a class of molecules, generally derived from vitamins, which function catalytically in enzyme systems.⁵ The acetic acid moiety which is bound by a high-energy bond (free energy 34.3 kJ/mole) to the -SH group of coenzyme A is a precursor to fatty acids, steroids, and other naturally occurring compounds, such as terpenes and acetogenins present in plants.^{6,7} The biosynthetic pathways for Acetyl CoA have been illustrated.⁶

In the transfer reaction by Acetyl CoA of the C_2 acetyl fragment, either the carboxyl group or the methyl group may react (electrophilic vs. nucleophilic reaction, respectively).⁷

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

Acetyl CoA, A2056, is soluble in water at 50 mg/mL. Aqueous solutions stored in aliquots at -20 °C are stable for no longer than 2 weeks.

Acetyl CoA is generally stable in neutral and moderately acidic solutions, even at elevated temperatures for a short time; aqueous solutions at pH 3.5–5 can be heated to 100 °C without decomposition.¹ Acetyl CoA hydrolyzes in strong acid and hydrolyzes more rapidly in alkaline solutions.

Storage/Stability

Store the product desiccated at -20 °C.

References

1

- Dawson, R.M.C. et al., Data for Biochemical Research, 3rd edition. Clarendon Press (Oxford, UK), pp. 116-117 (1993).
- Simon, E.J., and Shemin, D., J. Am. Chem. Soc., 75(10), 2520, (1953).
- 3. Stadtman, E.R., *Meth. Enzymol.*, **3**, 931-941 (1957).
- Bergmeyer, H.-U. et al., Methods of Enzymatic Analysis, 2nd edition, Vol. 4 (Bergmeyer, H.-U., ed.). Verlag Chemie/Academic Press, pp. 1988-2000 (1974).
- 5. Meyers, R.A., *Molecular Biology and Biotechnology: A Comprehensive Desk Reference*. VCH Publishers, Inc. (New York, NY), p. 960 (1995).
- 6. Lehninger, A.L., *Biochemistry*. Worth Publishers, Inc., p. 534 (1972).
- 7. Scott, T., and Eagleson, M., Concise Encyclopedia: Biochemistry, 2nd edition. Walter de Guyter Press, (New York, NY), p. 4 (1988).



Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose. The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.



