

For life science research only.
Not for use in diagnostic procedures.



cOmplete, Mini Protease Inhibitor Cocktail Tablets provided in *EASYpack*

 **Version: 06**

Content Version: March 2021

For the complete inhibition of proteases during extractions from animal and plant tissues or cells, yeast, and bacteria

Cat. No. 04 693 124 001 30 tablets
for 10 ml each

Store product at +2 to +8°C.

1.	General Information	3
1.1.	Contents	3
1.2.	Storage and Stability	3
	Storage Conditions (Product)	3
	Storage Conditions (Working Solution).....	3
1.3.	Additional Equipment and Reagent required	3
1.4.	Application	4
2.	How to Use this Product	5
2.1.	Before you Begin	5
	Safety Information	5
	Working Solution.....	5
	Preparation of 7x-concentrated stock solution	5
2.2.	Parameters	5
	Absorption.....	5
	Working Concentration.....	5
3.	Results	6
4.	Additional Information on this Product	6
4.1.	Quality Control.....	6
5.	Supplementary Information	7
5.1.	Conventions.....	7
5.2.	Changes to previous version	7
5.3.	Ordering Information.....	7
5.4.	Trademarks.....	8
5.5.	License Disclaimer	8
5.6.	Regulatory Disclaimer.....	8
5.7.	Safety Data Sheet.....	8
5.8.	Contact and Support.....	8

1. General Information

1.1. Contents

Package	Label	Function / Description	Content
1	cOplete, Mini, Protease inhibitor cocktail tablets	<ul style="list-style-type: none"> Individually packaged tablets. Each tablet is sufficient for a volume of 10 ml solution. 	30 foil blister packs, 1 tablet each

1.2. Storage and Stability

Storage Conditions (Product)

When stored at +2 to +8°C, the product is stable through the expiration date printed on the label.

Package	Label	Storage
1	cOplete, Mini, Protease inhibitor cocktail tablets	Store dry at +2 to +8°C.

Storage Conditions (Working Solution)

Store stock solution at +2 to +8°C for 1 to 2 weeks or –15 to –25°C for at least 12 weeks.

1.3. Additional Equipment and Reagent required

For preparation of stock solution

- Double-distilled water or 100 mM phosphate buffer, pH 7.0

For extraction in the acid pH range

- Pepstatin*

For determination of proteolytic activity

- Universal Protease Substrate, Casein, resorufin-labeled*

1.4. Application

cOplete, Mini Protease inhibitor cocktail tablets are used for the inhibition of a broad spectrum of serine, cysteine, and metalloproteases, as well as calpains. Due to the optimized composition of the tablets, they show excellent inhibition effects and are well suited for the protection of proteins isolated from:

- Mammalian tissue
- Bacteria
- Yeast
- Plants

cOplete, Mini contains both reversible and irreversible protease inhibitors, therefore, we recommend the addition of cOplete, Mini to all stock buffers and solutions normally protected with protease inhibitors and not only during the initial purification steps.

Since cOplete, Mini contains EDTA (3.7 mg/tablet yields a 1 mM solution of EDTA in 10 ml), the extraction buffer should not contain divalent cations such as Ca^{2+} , Mg^{2+} , or Mn^{2+} otherwise the inhibition of the metalloproteases might be incomplete. If the protein of interest will be purified by IMAC (Immobilized Metal Affinity Chromatography), for example, poly-His tagged recombinant proteins, EDTA must be eliminated, for example, by dialysis prior to the chromatography.

Alternatively, the product cOplete, Mini, EDTA-free* can be used. These tablets are identical to cOplete, Mini, with the only difference that no EDTA or other chelating agents are present.

2. How to Use this Product

2.1. Before you Begin

Safety Information

Do not eat the tablets.

Working Solution

⚠ Carefully push the tablet through the foil packaging using the base of your thumb (not fingernail) to prevent the breakage of tablets.

One cOplete, Mini Protease inhibitor tablet is sufficient for the inhibition of the proteolytic activity in 10 ml extraction solution. If very high proteolytic activity is present, use one tablet for 7 ml extraction buffer. Tablets can be added directly to the extraction medium. Alternatively, a 7x-concentrated stock solution can be prepared.

If it is necessary to inhibit proteolytic activity in a larger volume, use cOplete* (1 tablet for 50 ml extraction solution). The composition of the cOplete tablet is identical to the normal cOplete, Mini tablet, therefore comparable results are achieved with both products.

Preparation of 7x-concentrated stock solution

Dissolve one cOplete, Mini Protease inhibitor tablet in 1.5 ml double-distilled water or in 1.5 ml 100 mM phosphate buffer, pH 7.0.

2.2. Parameters

Absorption

A solution of 1 tablet cOplete, Mini inhibitor dissolved in 10 ml water has an absorption of 0.08 at 280 nm.

Working Concentration

1 tablet per 10 ml extraction solution.

3. Results

cOplete, Mini tablets efficiently inhibit serine, cysteine, and metalloproteases in a broad range.

Extract from	Type of protease present
Animal tissues	Serine, cysteine, and metalloproteases
Plants	Serine and cysteine proteases
Bacterial	Serine and metalloproteases

Occasionally, aspartic proteases (acid proteases) can interfere upon isolations from animal tissues. These proteases, however, exhibit pronounced activities only at low pH values. If extraction or single isolation steps must be performed at this pH range, add Pepstatin* to inhibit aspartic protease activity as a precaution.

Typical values for the inhibition of different proteases and protease mixtures by cOplete, Mini are shown in the following table. One cOplete, Mini tablet was added per 10 ml incubation solution. Proteolytic activity was determined with the Universal Protease Substrate, Casein, resorufin-labeled*. When extractions or single-step isolations are necessary in the acid pH range, include Pepstatin* along with cOplete, Mini tablets to ensure aspartic (acid) protease inhibition. All experiments were performed at +15 to +25°C.

Protease respective protease mixture	Enzyme concentration [mg/ml]	pH value	% Inhibition after immediate addition to the protease	% Inhibition after 1 hour incubation with the protease
Pancreas extract	0.02	7.8	85	96
Pronase	0.005	7.8	92	98
Thermolysin	0.0005	7.8	98	98
Chymotrypsin	0.0015	7.8	85	93
	0.003	7.8	90	93
Papain	0.33	7.8	66	77

The inhibitory power of cOplete, Mini has been demonstrated with many proteases and protease mixtures. In these experiments, drastically higher concentrations of proteases were used compared to the concentration usually present in extracts, see table above.

4. Additional Information on this Product



4.1. Quality Control

The inhibitory activity of each lot is tested with a concentrated pancreas extract and a concentrated pronase solution. Thereby, the proteolytic activities are typically inhibited by 95% after one hour (detection with Universal Protease Substrate Casein, resorufin-labeled*).

5. Supplementary Information

5.1. Conventions

To make information consistent and easier to read, the following text conventions and symbols are used in this document to highlight important information:

Text convention and symbols	
	<i>Information Note: Additional information about the current topic or procedure.</i>
	Important Note: Information critical to the success of the current procedure or use of the product.
① ② ③ etc.	Stages in a process that usually occur in the order listed.
① ② ③ etc.	Steps in a procedure that must be performed in the order listed.
* (Asterisk)	The Asterisk denotes a product available from Roche Diagnostics.

5.2. Changes to previous version

Layout changes.
Editorial changes.

5.3. Ordering Information

Product	Pack Size	Cat. No.
Reagents, kits		
cOplete	20 tablets in a glass vial, for 50 ml each	11 697 498 001
	3 x 20 tablets in glass vials, for 50 ml each	11 836 145 001
cOplete, Mini, EDTA-free	25 tablets in a glass vial, for 10 ml each	11 836 170 001
Universal Protease Substrate	40 mg	11 734 334 001
Pepstatin	2 mg	10 253 286 001
	10 mg	11 359 053 001
	50 mg	11 524 488 001

5. Supplementary Information

5.4. Trademarks

All product names and trademarks are the property of their respective owners.

5.5. License Disclaimer

For patent license limitations for individual products please refer to:

List of biochemical reagent products.

5.6. Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

5.7. Safety Data Sheet

Please follow the instructions in the Safety Data Sheet (SDS).

5.8. Contact and Support

To ask questions, solve problems, suggest enhancements or report new applications, please visit our **Online Technical Support Site.**

To call, write, fax, or email us, visit **sigma-aldrich.com**, and select your home country. Country-specific contact information will be displayed.

