

# Product Information

## CK1 $\gamma$ 3, active, GST-tagged, human PRECISIO® Kinase recombinant, expressed in Sf9 cells

Catalog Number **SRP5241**

Storage Temperature –70 °C

Synonym: CSNK1G3

### Product Description

CK1 $\gamma$ 3 is a member of the CK1 family of serine/threonine protein kinases, which plays an important role in diverse cell processes, including DNA replication and repair. CK1 $\gamma$ 3 is a ubiquitously expressed protein kinase found in the nuclei, cytoplasm, and membrane fractions of eukaryotic cells.<sup>1</sup> CK1 $\gamma$ 3 preferentially phosphorylates acidic substrates using ATP as a phosphate donor. The kinase domain of CK1 isoforms have been shown to associate with protein kinase C-potentiated inhibitor protein of 17 kDa, called CPI-17.<sup>2</sup> CPI-17 specifically inhibits myosin light chain phosphatase and this effect is potentiated when it is phosphorylated on Thr<sup>38</sup> by protein kinases such as the CK1 isoforms.

Full length recombinant human CK1 $\gamma$ 3 was expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag. The CK1 $\gamma$ 3 gene accession number is NM\_001031812. Recombinant protein stored in 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM glutathione, 0.1 mM EDTA, 0.25 mM DTT, 0.1 mM PMSF, and 25% glycerol.

Molecular mass: ~73 kDa.

Purity: 70–95% (SDS-PAGE, see Figure 1)

Specific Activity: 17–23 nmole/min/mg (see Figure 2)


### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.


### Storage/Stability

The product ships on dry ice and storage at –70 °C is recommended. After opening, aliquot into smaller quantities and store at –70 °C. Avoid repeated handling and multiple freeze/thaw cycles.

**Figure 1.**  
SDS-PAGE Gel of Typical Lot  
70–95% (densitometry)



**Figure 2.**  
Specific Activity of Typical Lot  
17–23 nmole/min/mg



### Procedure

#### Preparation Instructions

Kinase Assay Buffer – 25 mM MOPS, pH 7.2, 12.5 mM glycerol 2-phosphate, 25 mM MgCl<sub>2</sub>, 5 mM EGTA, and 2 mM EDTA. Just prior to use, add DTT to a final concentration of 0.25 mM.

Kinase Dilution Buffer – Dilute the Kinase Assay Buffer 5-fold with a 50 ng/ $\mu$ l BSA solution.

Kinase Solution – Dilute the active CK1 $\gamma$ 3 (0.1  $\mu$ g/ $\mu$ l) with Kinase Dilution Buffer to the desired concentration.

Note: The specific activity plot may be used as a guideline (see Figure 2). It is recommended the researcher perform a serial dilution of active CK1 $\gamma$ 3 kinase for optimal results.

10 mM ATP Stock Solution – Dissolve 55 mg of ATP in 10 ml of Kinase Assay Buffer. Store in 200  $\mu$ l aliquots at -20 °C.

$\gamma$ -<sup>33</sup>P-ATP Assay Cocktail (250  $\mu$ M) – Combine 5.75 ml of Kinase Assay Buffer, 150  $\mu$ l of 10 mM ATP Stock Solution, 100  $\mu$ l of  $\gamma$ -<sup>33</sup>P-ATP (1 mCi/100  $\mu$ l). Store in 1 ml aliquots at -20 °C.

Substrate Solution – Casein, Dephosphorylated, a protein substrate, was diluted in distilled water to a final concentration of 1 mg/ml.

1% phosphoric acid solution – Dilute 10 ml of concentrated phosphoric acid to a final volume of 1 L with water.

#### Kinase Assay

This assay involves the use of the <sup>33</sup>P radioisotope. All institutional guidelines regarding the use of radioisotopes should be followed.

1. Thaw the active CK1 $\gamma$ 3, Kinase Assay Buffer, Substrate Solution, and Kinase Dilution Buffer on ice. The  $\gamma$ -<sup>33</sup>P-ATP Assay Cocktail may be thawed at room temperature.
2. In a pre-cooled microcentrifuge tube, add the following solutions to a volume of 20  $\mu$ l:  
10  $\mu$ l of Kinase Solution  
5  $\mu$ l of Substrate Solution  
5  $\mu$ l of cold water (4 °C)
3. Set up a blank control as outlined in step 2, substituting 5  $\mu$ l of cold water (4 °C) for the Substrate Solution.
4. Initiate each reaction with the addition of 5  $\mu$ l of the  $\gamma$ -<sup>33</sup>P-ATP Assay Cocktail, bringing the final reaction volume to 25  $\mu$ l. Incubate the mixture in a water bath at 30 °C for 15 minutes.
5. After the 15 minute incubation, stop the reaction by spotting 20  $\mu$ l of the reaction mixture onto an individually pre-cut strip of phosphocellulose P81 paper.

6. Air dry the pre-cut P81 strip and sequentially wash in the 1% phosphoric acid solution with constant gentle stirring. It is recommended the strips be washed a total of 3 times of ~10 minutes each.
7. Set up a radioactive control to measure the total  $\gamma$ -<sup>33</sup>P-ATP counts introduced into the reaction. Spot 5  $\mu$ l of the  $\gamma$ -<sup>33</sup>P-ATP Assay Cocktail on a pre-cut P81 strip. Dry the sample for 2 minutes and read the counts. Do not wash this sample.
8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.
9. Determine the corrected cpm by subtracting the blank control value (see step 3) from each sample and calculate the kinase specific activity

#### Calculations:

1. Specific Radioactivity (SR) of ATP (cpm/nmole)

$$SR = \frac{\text{cpm of } 5 \mu\text{l of } \gamma\text{-}^{33}\text{P-ATP Assay Cocktail}}{\text{nmole of ATP}}$$

cpm – value from control (step 7)

nmole – 1.25 nmole (5  $\mu$ l of 250  $\mu$ M ATP Assay Cocktail)

2. Specific Kinase Activity (SA) (nmole/min/mg)

$$\text{nmole/min/mg} = \frac{\Delta \text{cpm} \times (25/20)}{SR \times E \times T}$$

SR = specific radioactivity of the ATP (cpm/nmole ATP)

$\Delta$ cpm = cpm of the sample – cpm of the blank (step 3)

25 = total reaction volume

20 = spot volume

T = reaction time (minutes)

E = amount of enzyme (mg)

#### References

1. Kusuda, J. et al., Cloning and chromosome mapping of the human casein kinase I  $\gamma$ -3 gene (CSNK1G3). *Cytogenet. Cell Genet.*, **83**, 101-103 (1998).
2. Zemlickova, E. et al., Association of CPI-17 with protein kinase C and casein kinase I. *Biochem Biophys Res Commun.*, **316(1)**, 39-47 (2004).

PRECISIO is a registered trademark of Sigma-Aldrich Co. LLC.

RC,MAM 10/11-1