

Product Information

Lipodisq® Styrene:Maleic Anhydride Copolymer

2:1, Pre-Hydrolyzed

SAE0062

Storage Temperature: Room Temperature

Product Description

Membrane proteins have historically proven to be a technically difficult class of proteins to study. This is primarily due to the difficulty of maintaining their solubility, stability, and function outside of their original hydrophobic and lipophilic contexts, in standard aqueous experimental conditions. Several experimental methodologies, such as Nanodisc technology, have been developed to enable the study of isolated membrane proteins in more conventional hydrophilic laboratory environments. Polymers of styrene and maleic acid (SMA) have emerged as an additional tool to facilitate the solubilization of membrane proteins.¹⁻⁴

Lipodisq® polymers contain styrene and maleic acid monomer units. Lipodisq® polymers may be used to form nano-sized discoidal particles that are suitable for incorporation of membrane proteins.^{5,6} Lipodisq® polymers can solubilize commonly used lipids, for example dimyristoylphosphatidylcholine (DMPC), without the use of detergents.⁷ In general, Lipodisq® polymers are potentially useful in mitigating the use of detergents for solubilization of membrane proteins.^{1,7-9}

This product is the pre-hydrolyzed form of the SMA polymer, with the styrene and maleic acid monomers present in the polymer in a 2:1 ratio. Of the four standard SMA ratios, 1:1, 2:1, 3:1, and 4:1, the SMA 2:1 polymer has been shown to have the broadest pH range for solubilization.¹⁰

Product Profile

Functional pH range: 5.5–8.0

M_w (weight average): ~ 7,500 DaM_n (number average): ~ 2,800 Da

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

References

1. Knowles, T.J. et al., *J. Am. Chem. Soc.*, **131(22)**: 7484-7485 (2009).
2. Dörr, J.M. et al., *Eur. Biophys. J.*, **45(1)**: 3-21 (2016).
3. Swainsbury, D.J.K. et al., *Biochim. Biophys. Acta*, **1859(10)**: 2133-2143 (2017).
4. Domínguez Pardo, J.J. et al., *Biophys. J.*, **115(1)**: 129-138 (2018).
5. Zhang, R. et al., *Biochim. Biophys. Acta.*, **1848(1 Pt B)**: 329-333 (2015).
6. Cuevas Arenas, R. et al., *Sci. Rep.*, **7**: 45875 (2017).
7. Orwick-Rydmark, M. et al., *Nano Lett.*, **12(9)**: 4687-4692 (2012).
8. Orwick, M.C. et al., *Angew. Chem. Int. Ed. Engl.*, **51(19)**: 4653-4657 (2012).
9. Dörr, J.M. et al., *Proc. Nat. Acad. Sci. USA*, **111(52)**: 18607-18612 (2014).
10. Scheidelaar, S. et al., *Biophys. J.*, **111(9)**: 1974-1986 (2016).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck operates as MilliporeSigma in the U.S. and Canada.

Merck and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.
SAE0062pis Rev 06/21

The Merck logo is displayed in a bold, red, sans-serif font.