

Product Information

Bone Morphogenetic Protein 2 Human

BMP-2, Recombinant, Expressed in HEK 293 Cells,
HumanKine®, Suitable for Cell Culture

H4791

Storage Temperature –20 °C

Synonym: BMP-2

Product Description

HumanKine® Bone Morphogenetic Protein 2 is produced from a DNA sequence encoding the human BMP-2 protein, expressed in HEK 293 cells. It is a glycosylated homodimer linked by a single disulfide bond with an apparent molecular mass of 30–38 kDa.

Bone Morphogenetic Protein 2 is a member of the TGF- β superfamily of cytokines that affect bone and cartilage formation. It is important for skeletal development during embryogenesis. BMP-2 induces chondrocyte formation, osteoblast differentiation, and is involved in embryo dorsal-ventral patterning and organogenesis.

It has been reported that BMP-2 inhibits estradiol-induced proliferation of human breast cancer cells.¹ BMP-2 signaling mediates apoptosis by activation of the TAK1-p38 kinase pathway that is negatively regulated by Smad6.² Cellular responses to BMP-2 are mediated by the formation of hetero-oligomeric complexes of type I and type II serine/threonine kinase receptors,³ which play significant roles in BMP binding and signaling. One BMP type II receptor and two BMP type I receptors have been identified. Both BMP type I receptors bind BMP-2 with high-affinity in the absence of BMP receptor type II.

This product is lyophilized from a solution of 50 mM sodium acetate, pH 4.5, with 50 mM NaCl.

EC50 60 ng/mL

The specific activity is determined by the dose dependent induction of alkaline phosphatase production in the ATDC-5 cell line (mouse chondrogenic cell line).

Purity \geq 95% (SDS-PAGE)

Endotoxin level \leq 1 EU/ μ g

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses.

Preparation Instructions

Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile PBS containing 0.1% endotoxin-free recombinant human serum albumin

Storage/Stability

Store the product at –20 °C. The lyophilized product remains active for one year at –20 °C.

Upon reconstitution, the cytokine can be stored at 2–8 °C for short term only, or at –20 °C to –80 °C in aliquots for long term. Avoid repeated freeze-thaw cycles.

References

1. Ghosh-Choudury, N., et al., *Bone morphogenetic protein-2 induces cyclin kinase inhibitor p21 and hypophosphorylation of retinoblastoma protein in estradiol-treated MCF-7 human breast cancer cells*. *Biochem. Biophys. Acta*, **497**: 186-196 (2000).
2. Kimura, N., et al., *BMP2-induced apoptosis is mediated by activation of the TAK1-p38 kinase pathway that is negatively regulated by Smad6*. *J. Biol. Chem.*, **275**: 17647-17652 (2000).
3. Kawabata, M., et al., *Signal transduction by bone morphogenetic proteins*. *Cytokine Growth Factor Rev.*, **9**: 49-61 (1998).
4. Chen, D. et al., *Growth Factors*, **22**: 233-241 (2004).
5. Nakamura, K. et al., *Exp. Cell research*, **250**: 351-363 (1999).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

H4791pis Rev 06/21

**MILLIPORE
SIGMA**