

3050 Spruce Street, St. Louis, MO 63103 USA
Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757
email: techservice@sial.com sigma-aldrich.com

# **Product Information**

SUMO-1, human, GST-tagged recombinant, expressed in *E. coli* 

Catalog Number **S0439** Storage Temperature –70 °C

Synonyms: Sentrin-1, GMP-1, PIC1

## **Product Description**

The class of post-translational protein modifications (PTM) known as **s**mall **u**biquitin-related **mo**difiers (SUMO) are part of the PTM class of ubiquitin and/or ubiquitin-like proteins (UBL). SUMO-1 is one of 5 SUMO paralogs expressed in mammalian cells. SUMO-1 is a protein of 101 amino acids with a molecular mass of ~11.5 kDa. SUMO-1 has a low but significant homology with ubiquitin.

Almost all the SUMO-1 in cells is conjugated to other proteins. Very little free SUMO-1 has been found. The proteins to which SUMO-1 is conjugated include p53 (tumor suppressor protein),  $I\kappa B\alpha$ , and Ran GTPase-activating protein 1 (RanGAP1). Proteins that cannot be ubiquinated and degraded in that pathway are processed by conjugation to SUMO-1. The factors required for conjugation have been elucidated with RanGAP1. Conjugation proceeds without the equivalent of an  $E_3$  ubiquitin ligase. In the presence of SUMO-1 activating enzyme (SAE1/SAE2), Ucbh9, and ATP, SUMO-1 is efficiently conjugated to  $I\kappa B\alpha$ . SUMO-1 labeling/modification is implicated in the progression of  $G_2$  to M phase cell cycle progression.

This SUMO-1-GST fusion protein is produced from a DNA sequence corresponding to human SUMO-1 fused to a GST-tag, and is expressed in *E. coli* cells. The molecular mass of SUMO-1-GST is ~38.5 kDa. This SUMO-1-GST product is supplied as a solution in 50 mM HEPES, pH 8.0, containing 150 mM NaCl and 1 mM DTT.

## **Precautions and Disclaimer**

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

### Storage/Stability

The product ships on dry ice, and storage at -70 °C is recommended. The protein is stable to multiple freeze/thaw cycles.

#### References

- Hickey, C.M. et al., Nat. Rev. Mol. Cell Biol., 13(12), 755-766 (2012).
- 2. Pichler, A. et al., Biomol. Concepts, **8(1)**, 13-36 (2017).
- 3. Mahajan, R. et al., Cell, 88(1), 97-107 (1997).
- Desterro, J.M.P. et al., Mol. Cell, 2(2), 233-239 (1998).
- 5. Rodriguez, M.S. *et al.*, *EMBO J.*, **18(22)**, 6455-6461 (1999).
- 6. Saitoh, H. et al., Curr. Biol., 8(2), 121-124 (1998).
- Desterro, J.M.P. et al., J. Biol. Chem., 274(15), 10618-10624 (1999).

RBG,FEB,JWM,GCY,MAM 07/18-1