

Product Information

Monoclonal Anti-Histone Deacetylase 11 (HDAC11)

Clone HDAC11-31
Purified Mouse Immunoglobulin

Product Number **H 2913**

Product Description

Monoclonal Anti-Histone Deacetylase 11 (HDAC11) (mouse IgG2b isotype) is derived from the HDAC11-31 hybridoma produced by the fusion of mouse myeloma cells (NS1) and splenocytes from BALB/c mice immunized with a synthetic peptide corresponding to amino acids 2-16 of human HDAC11, conjugating to KLH. The HDAC11 sequence in mouse and rat differs by three amino acids. The isotype is determined using Sigma ImmunoType™ Kit (Product Code ISO-1) and by a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents (Product Code ISO-2).

Monoclonal Anti-Histone Deacetylase 11 (HDAC11) recognizes human HDAC11 (approx. 39 kDa). The antibody epitope resides within amino acids 2-16 of human HDAC11. The antibody may be used in ELISA and immunoblotting.

The basic repeating unit of chromatin is the nucleosome, which is composed of a protein octamer containing two each of the core histones H2A, H2B, H3, and H4 surrounded by approximately 146 base pairs of DNA. Reversible acetylation of highly conserved lysine residues in the N-terminal tail domains of core histones plays an important role in transcriptional regulation, cell cycle progression, and development events. Several histone acetyltransferases (HATs) catalyze this acetylation reaction (e.g. GCN5, PCAF, p300/CBP, TAFII250, P/CAF, SRC-1, BRCA-2). Acetylation of the core histones is generally considered to be associated with gene activation, probably through maintenance of the unfolded structure of transcribing nucleosomes.^{1,2}

Histone acetylation is a dynamic process where levels are determined by the net activities of HATs and the competing enzymes histone deacetylases (HDACs).³ Both activities are associated with the nuclear matrix. Eleven different mammalian HDACs have been described. HDAC1-3 and 8 (Class I) are similar to yeast Rpd3 protein, while HDAC4-7, 9 and 10 (Class II) are similar to yeast Hda1 protein.^{4, 5, 13} The activities of the histone deacetylases are often, but not always, associated with transcriptional repression and nucleosome condensation.^{6, 7} HDAC1, HDAC2, and several others are the catalytic subunits of different multiprotein regulatory complexes.⁸ Other components of such complexes may include corepressors such as mSin3, N-CoR, SMRT, and associated proteins such as SAP18, SAP30, RbAp46, RbAp48, and c-Ski oncogenic protein, a protein involved in DNA methylation. Nucleosome remodeling and deacetylation (NRD) complexes containing HDAC1, HDAC2, Mi-2 (CH3, CH4) dermatomyositis specific autoantigen, and MAT2 (metastasis-associated protein) protein, have been described. It is therefore assumed that ATP-dependent nucleosome remodeling activity and histone deacetylation may be interconnected or interdependent.^{9, 10} Recruitment of the multiprotein complexes to promoter sites occurs by many sequence specific DNA-binding proteins such as unliganded nuclear hormone receptors, DP1-E2F, YY1, and Rb family of transcription factors, transcriptional repressors, and tumor suppressors (e.g. BRCA1). Aberrant recruitment of HDACs by certain oncoproteins may occur in certain neoplastic diseases.¹¹

HDAC11 is a new member of the HDAC family. The protein does not belong to HDAC class I or II but is related to a common ancestral gene(s) from which the eukaryotic HDACs evolved.¹² HDAC11 has a molecular weight of 39 kDa and is mainly expressed in brain, heart, skeletal muscle, kidney, testis, and cancer cells. The protein consists of one catalytic domain and is found in a complex with HDAC6.¹²

Reagent

Monoclonal Anti-Histone Deacetylase 11 (HDAC11) is supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Antibody Concentration: Approx. 2 mg/ml.

Precautions and Disclaimer

Due to the sodium azide content, a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing is not recommended. Storage in frost-free freezers is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

By immunoblotting, a working antibody concentration of 2-4 µg/ml is recommended using HeLa cell extracts.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilution by titration.

References

1. Bauer, W.R., et al., *J. Mol. Biol.*, **236**, 685-690 (1994).
2. Lee, D.Y., et al., *Cell*, **72**, 73-84 (1993).
3. Taunton, J., et al., *Science*, **272**, 408-411 (1996).
4. Grozinger, C.M., et al., *Proc. Natl. Acad. Sci. USA*, **96**, 4868-4873 (1999).
5. Fischle, W., et al., *J. Biol. Chem.*, **274**, 11713-11720 (1999).
6. Wolffe, A.P., *Science*, **272**, 371-372 (1996).
7. Grunstein, M., *Nature*, **389**, 349-352 (1997).
8. Hassig, C.A., et al., *Proc. Natl. Acad. Sci. USA*, **95**, 3519-3524 (1998).
9. Tong, J.K., et al., *Nature*, **395**, 917-921 (1998).
10. Zhang, Y., et al., *Cell*, **95**, 279-289 (1998).
11. Lin, R.J. et al., *Nature*, **391**, 811-814 (1998).
12. Gao, L., et al., *J. Biol. Chem.*, **277**, 25748-25755 (2002).

KAA/EK 02/04

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.