

3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

ProductInformation

N-Methyl-D-glucamine

Product Number M 2004
Store at room temperature
Replacement for Product Code M4,700-0

Product Description

Molecular Formula: $C_7H_{17}NO_5$ Molecular Weight: 195.2 CAS Number: 6284-40-8 Melting point: 128-129 °C Specific rotation: -15.7 to -17.3° (10 mg/ml in water at 25 °C)

pK_a: 9.5 (20 °C)¹

Synonyms: NMDG, meglumine

This product is used as a Na⁺ substitute in systems to maintain physiological solutions as isotonic, but without sodium. It has been used, among other things, in studying cells from rabbit cerebral artery,² rabbit cardiacmyocytes,³ perfused rat intestine,⁴ and frog skeletal muscle.⁵

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

Preparation Instructions

This product is soluble in water (50 mg/ml). It is reported to be soluble up to 1g/ml in water at 25 °C. ⁶ A 1% aqueous solution has a pH of approximately 10.5.

References

- Martindale The Extra Pharmacopoeia, 29th ed., Reynolds, J. E. F., ed., The Pharmaceutical Press (London, England: 1989), p. XXVI.
- Kim, J. K., et al., Influence of Extracellular Na⁺ Removal on Cytosolic Ca²⁺ Concentration in Smooth Muscle Cells of Rabbit Cerebral Artery. J. Smooth Muscle Res., 35, 135-145 (1999).
- Clemo, H. F., and Baumgarten, C. M., Atrial Natriuretic Factor Decreases Cell Volume of Rabbit Atrial and Ventricular Myocytes. Am. J. Physiol., 260(4 Pt 1), C681-C690 (1991).
- Wapnir, R. A., and Stiel, L., Intestinal Absorption of Copper: Effect of Sodium. Proc. Soc. Exp. Biol. Med., 185(3), 277-282 (1987).
- Abercrombie, R. F., et al., The Intracellular pH of Frog Skeletal Muscle: Its Regulation in Isotonic Solutions. J. Physiol., 345, 175-187 (1983).
- The Merck Index, 11th ed., Entry# 5995.

CMH/RXR 6/03