



MOUSE ANTI-NERVE GROWTH FACTOR-RECEPTOR, p75 MONOCLONAL ANTIBODY

CATALOG NUMBER:	MAB5386	QUANTITY:	100 µg
LOT NUMBER:			
ALTERNATE NAMES:	low affinity NGF-Receptor; p75, NGFR	EPITOPE:	within aa 1-160, extracellular domain of human p75.
CLONE NAME:	ME20.4	HOST/ISOTYPE:	Ms IgG1
BACKGROUND:	The low affinity NGFR (Nerve growth factor receptor) is a 75kDa membrane-spanning glycoprotein lacking intrinsic tyrosine kinase activity. p75NGFR interacts with TrkA, the high affinity NGF receptor and potentiates TrkA signaling at low NGF concentrations. The p75 receptor binds nerve growth factor, brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4 with varying specificities. The p75NGFR plays an important role in neurotrophic factor signaling and has been shown to modulate the susceptibility of selective cellular populations to programmed cell death. It is expressed on many neuronal cells types including many embryonic forms and the receptor can be used to isolate neuronal progenitor cells		
SPECIFICITY:	Reacts with the low affinity nerve growth factor receptor (NGFR p75). The epitope recognized by the antibody maps within the third or fourth cysteine-rich repeats of the NGFR molecule. Reactivity has not been observed with rat, chicken, or mouse NGFR.		
IMMUNOGEN:	Human melanoma cell line (Ross, 1984).		
APPLICATIONS:	Western blot: RIPA lysates of human HS294T cells; 1 µg/mL; recognizes ~69-75kDa band depending upon gel conditions. Immunohistochemistry: 5-10 µg/mL on frozen tissue. Immunohistochemistry on formalin-fixed, paraffin embedded sections. Immunocytochemistry: 5-10 µg/mL EM Immunohistochemistry {Caneva, et al 1995}. http://www.scripps.edu/bcmd/pdfarea/issue_09/soligo.pdf Immunoabsorptive purification of NGF-receptors. Inhibition of NGF binding to p-75NGFR Immunoprecipitation of p75 FACS: unfixed cells RIA Optimal working dilutions must be determined by the end user.		
SPECIES REACTIVITY:	Reacts with Primate, Rabbit, Porcine (Pig), Human, Feline (Cat), Canine (Dog), Raccoon and Sheep (Ovine). Does not react with Rat, Mouse and Chicken.		
FORMAT:	Purified immunoglobulin.		
PRESENTATION:	Lyophilized. Reconstitute to 1 mg/mL with sterile distilled water.		



STORAGE/HANDLING: Maintain for 1 year at 2–8°C from date of shipment. Aliquot to avoid repeated freezing and thawing. For maximum recovery of product, centrifuge the original vial after reconstitution and prior to removing the cap.

REFERENCES: Tremere, LA, et al. (2000). Antibody for human p75 LNTR identifies cholinergic basal forebrain of non-primate species. *Neuroreport* 11:2177-2183.

Ross, AH, et al (1984). Characterization of nerve growth factor receptor in neural crest tumors using monoclonal antibodies. *PNAS.USA* 81:6681-6685.

Important Note: *During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200 μ L or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.*

FOR RESEARCH USE ONLY; NOT FOR USE IN DIAGNOSTIC
PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION

Unless otherwise stated in our catalog or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.