

Product Information

Anti-ARG1 antibody, Mouse monoclonal
clone ARG1-6, purified from hybridoma cell culture

Catalog Number **SAB4200509**

Product Description

Anti-ARG1 (mouse IgG2b isotype) is derived from the hybridoma ARG1-6 produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with a synthetic peptide corresponding to an internal region of human ARG1 (GenElD: 383), conjugated to KLH. The corresponding sequence differs by 3 amino acids in rat and by 4 amino acids in mouse ARG1. The isotype is determined by ELISA using Mouse Monoclonal Antibody Isotyping Reagents, Catalog Number ISO2. The antibody is purified from culture supernatant of hybridoma cells grown in a bioreactor.

Anti- ARG1, clone ARG1-6 recognizes human ARG1. The antibody may be used in various immunochemical techniques including immunoblotting (~40 kDa) and immunoprecipitation. Detection of the ARG1 band by immunoblotting is specifically inhibited by the immunizing peptide

Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exist (types I and II), which differ in their tissue distribution, subcellular localization, immunologic cross reactivity and physiologic function. Arginases have been implicated in many disease processes, including vascular diseases, pulmonary diseases, infectious diseases, immune cell dysfunction and cancer. The type I isoform encoded by the ARG1 gene, is a cytosolic enzyme expressed predominantly in the liver as a component of the urea cycle. Inherited deficiency of this enzyme results in argininemia, an autosomal recessive disorder characterized by hyperammonemia. Two transcript variants encoding different isoforms have been found for this gene.¹⁻⁵

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody Concentration: ~ 1.0 mg/mL

©2021 Sigma-Aldrich Co. LLC. All rights reserved. SIGMA-ALDRICH is a trademark of Sigma-Aldrich Co. LLC, registered in the US and other countries. Sigma brand products are sold through Sigma-Aldrich, Inc. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see product information on the Sigma-Aldrich website at www.sigmaaldrich.com and/or on the reverse side of the invoice or packing slip.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze at -20 °C in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working concentration of 0.25-0.50 µg/mL is recommended using whole extracts of HEK-293T cells overexpressing human ARG1.

Immunoprecipitation: a working amount of 2.5-5.0 µg is recommended using lysates of HEK-293T cells overexpressing human ARG1.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.

References

1. Morris, S.M.Jr., *Br. J. Pharmacol.*, **157**, 922-930 (2009).
2. Jenkinson, C.P., et al., *Comp. Biochem. Physiol. B Biochem. Mol. Biol.*, **114**, 107-132 (1996).
3. Di Costanzo, L., et al., *Proc. Natl. Acad. Sci. USA*, **102**, 13058-13063 (2005).
4. Uchino, T., et al., *Hum. Genet.*, **96**, 255-260 (1995).
5. Morris, S.M.Jr., et al., *Curr. Opin. Clin. Nutr. Metab. Care*, **15**, 64-70 (2012).

RC,ST,KCP,PHC 04/21-1